arXiv:1512.03786v1 [math.RT] 28 Nov 2015

MATRICIAL REPRESENTATIONS OF CERTAIN FINITELY PRESENTED GROUPS GENERATED BY ORDER-2 GENERATORS AND THEIR APPLICATIONS

RYAN GOLDEN AND ILWOO CHO

Abstract. In this paper, we study matricial representations of certain finitely presented groups Γ_N^2 with N-generators of order-2. As an application, we consider a group algebra $\mathcal{A}_2$ of Γ_2^2, under our representations. Specifically, we characterize the inverses g^{-1} of all group elements g in Γ_2^2, in terms of matrices in the group algebra $\mathcal{A}_2$. From the study of this characterization, we realize there are close relations between the trace of the radial operator of $\mathcal{A}_2$, and the Lucas numbers appearing in the Lucas triangle.

1. Introduction

In this paper, we consider certain *finite-dimensional representations* $(\mathbb{C}^n,\ \alpha^n)$ of a *finitely presented group* Γ_N^2,

(1.0.1)

$$\Gamma_N^2 \overset{def}{=} \left\langle \{x_1,\ ...,\ x_N\},\ \{x_1^2 = ... = x_N^2\} \right\rangle,$$

where the generators x_1, ..., x_N are *noncommutative indeterminants*. i.e., an algebraic structure Γ_N^2 is the *group* generated by N-*generators* x_1, ..., x_N equipped with its "noncommutative" binary operation $(\cdot)$, satisfying

$$x_1^2 = x_2^2 = ... = x_N^2 = e_N,$$

where e_N is the *group-identity of* Γ_N^2 such that:

$$ge_N = g = e_N g, \text{ for all } g \in \Gamma_N^2,$$

for all $N \in \mathbb{N}$. So, one may understand a group Γ_N^2 as the *quotient group*,

(1.0.2)

$$\Gamma_N^2 \overset{\text{Group}}{=} \mathcal{F}_N \ / \ \{g_j^2 = e : g_j \text{ are generators of } \mathcal{F}_N\},$$

where $\mathcal{F}_N$ is the *noncommutative* (*non-reduced*) *free group* $\left\langle \{g_j\}_{j=1}^N \right\rangle$ *with* N-*generators* g_1, ..., g_N, where e is the group-identity of the free group $\mathcal{F}_N$, and "$\overset{\text{Group}}{=}$" means "being group-isomorphic" (e.g., [1] and [2]).

By construction, all elements g of the group Γ_N^2 are of the form,

(1.0.3)

$$g = x_{i_1}^{k_1}\ x_{i_2}^{k_2}\ ...\ x_{i_n}^{k_n},$$

for some

$$i_1, ..., i_n \in \{1, ..., N\}, \text{ and } k_1, ..., k_n \in \mathbb{Z},$$

1991 *Mathematics Subject Classification.* 05A19, 30E50, 37E99, 44A60.

Key words and phrases. Finitely Presented Groups, Generators of Order 2, Representations, Matrix Groups, the Lucas Triangle, Lucas Numbers.

as non-reduced words in $\{x_1, ..., x_N\}$, where x_j^{-1} means the *group-inverse of* x_j, and hence, x_j^{-k} means $\left(x_j^{-1}\right)^k$, for all $k \in \mathbb{N}$, for all $j = 1, ..., N$.

In this paper, for an arbitrarily fixed $N \in \mathbb{N}$, we establish and study certain n-dimensional Hilbert-space representations $(\mathbb{C}^n, \alpha^n)$ of the groups Γ_N^2, for all $n \in \mathbb{N} \setminus \{1\}$. Under our representations, each element g of Γ_N^2 is understood as a matrix A_g acting on $\mathbb{C}^n$, for all $n \in \mathbb{N}$. Moreover, if

$$g = x_{i_1}^{k_1}...x_{i_l}^{k_l} \text{ as in (1.0.3)}$$

(as a "non-reduced" word in Γ_N^2), then there exists corresponding matrices A_{i_1}, ..., A_{i_l}, such that

$$A_g = A_{i_1}^{k_1}...A_{i_l}^{k_l} \text{ in } M_n(\mathbb{C})$$

(as a "reduced" word in $M_n(\mathbb{C})$), where $M_n(\mathbb{C})$ is the *matricial algebra* consisting of all $(n \times n)$-matrices, for all $n \in \mathbb{N}$.

1.1. **Motivation.** The readers may skip reading this sub-section. Here, we simply want to emphasize why we are interested in the groups Γ_N^2, for $N \in \mathbb{N}$.

Free probability is a branch of *operator algebra*. By considering free-distributional data, one can establish operator-valued *noncommutative* probability theory on (topological, or pure-algebraic) *algebras* as in classical probability theory (e.g., [3], [4] and cited papers therein). Also, such free-probabilistic data let us have structure theorems of given algebras under *free product* determined by given *linear functionals*. Here, the *independence* of classical probability is replaced by the so-called *freeness*. There are two approaches in free probability theory: *Voiculescu*'s original analytic approach (e.g., [4]), and *Speicher*'s combinatorial approach (e.g., [3]).

The second-named author, Cho, has considered connections between operator algebra theory, and Hecke-algebraic number theory via free probability recently, to provide tools for studying number-theoretic results with operator-algebraic techniques, and vice versa, by establishing certain representational and operator-theoretic models from combinatorial free probability settings of [3].

Especially, in the series of research papers [5], [2] and [1], Cho considered the *free-probabilistic representations* of the *Hecke algebras* $\mathcal{H}(G_p)$ generated by the *generalized linear* ((2×2)-*matricial*) *groups* $G_p = GL_2(\mathbb{Q}_p)$ over the *p-adic number fields* $\mathbb{Q}_p$, for *primes* p.

In the frontier paper [5], Cho and Gillespie established free probability models of certain subalgebras $\mathcal{H}_{Y_p}$ of the Hecke algebras $\mathcal{H}(G_p)$ by defining suitable linear functionals on $\mathcal{H}(G_p)$. In particular, they constructed free-probabilistic structures preserving number-theoretic data from $\mathcal{H}_{Y_p}$.

In [2], Cho extended free-probabilistic models of $\mathcal{H}_{Y_p}$ in the sense of [6] to those of $\mathcal{H}(G_p)$ fully, for all primes p. On such models, the *C^*-algebras* $\mathbb{H}(G_p)$ were constructed by realizing elements of $\mathcal{H}(G_p)$ as operators under free-probabilistic representations of $\mathcal{H}(G_p)$. And the operator-theoretic properties of *generating operators of* $\mathbb{H}(G_p)$ were studied there.

In [1], by studying certain types of *partial isometries* of $\mathbb{H}(G_p)$, Cho obtained the embedded non-abelian multiplicative groups G_N of $\mathbb{H}(G_p)$ generated by N-many partial isometries in the sense of [1]. In particular, it was shown there that:

(1.1.1)

$$G_N \overset{\text{Group}}{=} \Gamma_N^2, \text{ for all } N \in \mathbb{N},$$

where Γ_N^2 is in the sense of (1.0.1), satisfying (1.0.2).

To study detailed algebraic properties of G_N in $\mathcal{H}(G_p)$, and to investigate operator-algebraic properties of $C^*(G_N)$ in $\mathbb{H}(G_p)$, he used the isomorphic group Γ_N^2 of (1.0.1), and the corresponding group C^*-algebra $C^*\left(\Gamma_N^2\right)$.

Thus, for authors, it is natural to be interested in the groups Γ_N^2. In this paper, we study the groups Γ_N^2 of (1.0.1) pure-algebraically (independent from those in [1], [2] and [5]).

1.2. **Overview.** In this paper, we concentrate on studying the groups Γ_N^2 of (1.0.1) independently. The main purpose is to establish suitable *Hilbert-space representations* (other than those of [1]) of Γ_N^2. Fundamentally, we construct "finite-dimensional" Hilbert-space representations $(\mathbb{C}^n,\ \alpha^n)$ of Γ_N^2, for "all" $n \in \mathbb{N} \setminus \{1\}$.

Under our representation $(\mathbb{C}^n,\ \alpha^n)$ for $n \in \mathbb{N}$, each group element $g \in \Gamma_N^2$ is understood as a matrix A_g in the matricial algebra $M_n(\mathbb{C})$, consisting of all $(n \times n)$-matirices. To study an algebraic object $g \in \Gamma_N^2$, we will investigate functional properties of the corresponding matrices $A_g \in M_n(\mathbb{C})$ for g, for all $n \in \mathbb{N}$ (See Section 2).

As application, we consider a group algebra $\mathcal{A}_2$ of Γ_2^2, under our representations. Specifically, we characterize the inverses g^{-1} of all group elements g in Γ_2^2, in terms of matrices in the group algebra $\mathcal{A}_2$ generated by Γ_2^2 (See Section 3).

From the study of this characterization of Section 3, we show that there are close relations between the *trace* of the powers T^n of the *radial operator* T of $\mathcal{A}_2$, and the *Lucas numbers* in the *Lucas triangle* (See Section 4).

2. Finite-Dimensional Representations of Γ_N^2

In this section, we establish finite-dimensional Hilbert-space representations of the finitely presented groups Γ_N^2 with N-generators of order-2 in the sense of (1.0.1). i.e.,

$$\Gamma_N^2 = \left\langle \{x_j\}_{j=1}^N,\ \{x_j^2 = e_N\}_{j=1}^N \right\rangle,$$

where x_1, ..., x_N are noncommutative indeterminants (as generators), and e_N are the group-identities of Γ_N^2, for all $N \in \mathbb{N}$.

Fix $N \in \mathbb{N}$ throughout this section and the corresponding group Γ_N^2.

2.1. **A Matrix Group $\mathfrak{M}_2^2(N)$.** Let $M_2(\mathbb{C})$ be the (2×2)-matricial algebra, and let

$$\mathbb{C}^\times = \mathbb{C} \setminus \{0\} \text{ in } \mathbb{C}.$$

Suppose $\mathbb{C} \times \mathbb{C}^\times$ is the *Cartesian product* of $\mathbb{C}$ and $\mathbb{C}^\times$. Define now a function

$$A_2 : \mathbb{C} \times \mathbb{C}^\times \to M_2(\mathbb{C})$$

by

(2.1.1)

$$A_2(a,\, b) = \begin{pmatrix} a & b \\ \frac{1-a^2}{b} & -a \end{pmatrix},$$

for all $(a,\, b) \in \mathbb{C} \times \mathbb{C}^\times$. It is not difficult to check that:

Lemma 2.1. *Let A be the map in the sense of (2.1.1). Then*

(2.1.2)

$$(A_2(a,b))^2 = I_2, \text{ the identity matrix of } M_2(\mathbb{C}),$$

for all $(a,\, b) \in \mathbb{C} \times \mathbb{C}^\times$.

Proof. The proof of (2.1.2) is from the straightforward computation. ∎

It is also easy to verify that:

$$A_2\,(a_1, b_1)\ A_2(a_2,\, b_2) = \begin{pmatrix} \frac{a_1a_2b_2+b_1-b_1a_2^2}{b_2} & a_1b_2 - b_1a_2 \\ \frac{a_2b_2-a_2b_2a_1^2-a_1b_1+a_1b_1a_2^2}{b_1b_2} & \frac{b_2-b_2a_1^2+a_1a_2b_1}{b_1} \end{pmatrix},$$

and

$$A_2\,(a_2, b_2)\ A_2(a_1,\, b_1) = \begin{pmatrix} \frac{b_2-b_2a_1^2+a_1a_2b_1}{b_1} & b_1a_2 - a_1b_2 \\ \frac{-a_2b_2+a_2b_2a_1^2+a_1b_1-a_1b_1a_2^2}{b_1b_2} & \frac{a_1a_2b_2+b_1-b_1a_2^2}{b_2} \end{pmatrix},$$

and hence,

(2.1.3)

$$A_2\,(a_1, b_1)\ A_2(a_2, b_2) \neq A_2(a_2, b_2)\ A_2(a_1, b_1),$$

in general, for $(a_k,\, b_k) \in \mathbb{C} \times \mathbb{C}^\times$, for $k = 1,\, 2$. In particular, whenever

$$(a_k,\, b_k) \in \mathbb{C}^\times \times \mathbb{C}^\times \text{ in } \mathbb{C} \times \mathbb{C}^\times,$$

for $k = 1,\, 2$, and the pairs $(a_1,\, b_1)$ and $(a_2,\, b_2)$ are "distinct" in $\mathbb{C}^\times \times \mathbb{C}^\times$, the above noncommutativity (2.1.3) always holds.

Therefore, the images

$$\{A_2(a,\, b) : (a,\, b) \in \mathbb{C}^\times \times \mathbb{C}^\times\}$$

forms the noncommutative family in $M_2(\mathbb{C})$, by (2.1.3).

Now, take distinct pairs,

(2.1.3)′

$$(a_k,\, b_k) \in \mathbb{C}^\times \times \mathbb{C}^\times, \text{ for } k = 1,\, ...,\, N.$$

For the above chosen N-pairs $\{(a_k,\, b_k)\}_{k=1}^N$ of (2.1.3)′, one can construct the corresponding matrices,

(2.1.4)

$$A_2(a_k,\, b_k) \text{ in } M_2(\mathbb{C}), \text{ for } k = 1,\, ...,\, N,$$

by (2.1.1). Under the matrix multiplication on $M_2(\mathbb{C})$, let's construct the multiplicative subgroup $\mathfrak{M}_2^2(N)$ of $M_2(\mathbb{C})$ as

(2.1.5)

$$\mathfrak{M}_2^2(N) = \left\langle \{A_2(a_k,\ b_k)\}_{k=1}^N \right\rangle$$

generated by $A_2(a_k,\, b_k)$, for all $k = 1,\, ...,\, N$. i.e., all elements of $\mathfrak{M}_2^2(N)$ are the (2×2)-matrices in $M_2(\mathbb{C})$, satisfying

(2.1.6)

$$\left(A_2(a_k, b_k)\right)^2 = I_2 \text{ in } \mathfrak{M}_2^2(N),$$

for all $k = 1,\, ...,\, N$, be (2.1.2).

Definition 2.1. *Let $\mathfrak{M}_2^2(N)$ be the multiplicative subgroup (2.1.5) of $M_2(\mathbb{C})$. We call $\mathfrak{M}_2^2(N)$, the 2-dimensional, order-2, N-generator (sub-)group (of $M_2(\mathbb{C})$).*

One obtains the following algebraic characterization.

Theorem 2.2. *Let $\mathfrak{M}_2^2(N)$ be a 2-dimensional, order-2, N-generator group in $M_2(\mathbb{C})$. Then the groups $\mathfrak{M}_2^2(N)$ and the group Γ_N^2 of (1.0.1) are isomorphic. i.e.,*

(2.1.7)

$$\mathfrak{M}_2^2(N) \overset{Group}{=} \Gamma_N^2.$$

Proof. Let $\mathfrak{M}^2_2(N)$ be a 2-dimensional, order-2, N-generator group in $M_2(\mathbb{C})$. Then, by (2.1.6), each generator $A_2(a_k,\ b_k)$ satisfies

$$\left(A_2(a_k,\ b_k)\right)^2 = I_2, \text{ for all } k = 1,\ ...,\ N.$$

i.e., each generator $A_2(a_k,\ b_k)$ has order-2 indeed in $\mathfrak{M}^2_2(N)$.

Since $(a_k,\ b_k)$ are taken from $\mathbb{C}^\times \times \mathbb{C}^\times$, the generators $\{A_2(a_k,\ b_k)\}^N_{k=1}$ forms a noncommutative family in $M_2(\mathbb{C})$, by (2.1.3).

Observe now that there does not exist n-tuple $(j_1,\ ...,\ j_n)$ of "distinct" elements $j_1,\ ...,\ j_n$ in $\{1,\ ...,\ N\}$, for all $n \in \mathbb{N}$, such that

$$\prod_{l=1}^{n} A_2\left(a_{k_{j_l}},\ b_{k_{j_l}}\right) = A_2\left(a_{k_{j_0}},\ b_{k_{j_0}}\right), \text{ or } I_2,$$

for some $j_0 \in \{1,\ ...,\ N\}$, by the very definition (2.1.1) and (2.1.3).

These observations show that

$\mathfrak{M}^2_2(N) \overset{\text{Group}}{=} \mathcal{F}_N\ /\ \mathcal{R}_N$,

where $\mathcal{F}_N$ is the noncommutative free group $\left\langle \{g_j\}^N_{j=1} \right\rangle$ with N-generators $g_1,\ ...,\ g_N$, and $\mathcal{R}_N$ means the relator set,

$$\mathcal{R}_N = \{g_j^2 = e\}^N_{j=1},$$

where e is the group-identity of $\mathcal{F}_N$.

Recall that the group Γ^2_N of (1.0.1) is group-isomorphic to the quotient group $\mathcal{F}_N\ /\ \mathcal{R}_N$, by (1.0.2). Therefore, one has that

$$\mathfrak{M}^2_2(N) \overset{\text{Group}}{=} \mathcal{F}_N\ /\ \mathcal{R}_N \overset{\text{Group}}{=} \Gamma^2_N.$$

Therefore, the 2-dimensional, order-2, N-generator group $\mathfrak{M}^2_2(N)$ is group-isomorphic to the group Γ^2_N of (1.0.1). ∎

By (2.1.7), two groups $\mathfrak{M}^2_2(N)$ and Γ^2_N are isomorphic from each other, equivalently, there does exist a group-isomorphism,

$$\alpha^2 : \Gamma^2_N \to \mathfrak{M}^2_2(N),$$

satisfying

(2.1.8)

$$\alpha^2\left(x_j\right) = A_2\left(a_j,\ b_j\right), \text{ for all } j = 1,\ ...,\ N.$$

The above structure theorem (2.1.7), equivalently, the existence of the group-isomorphism α^2 of (2.1.8) provides a 2-dimensional Hilbert-space representation $\left(\mathbb{C}^2,\ \alpha^2\right)$ of Γ^2_N.

Theorem 2.3. *There exists the 2-dimensional Hilbert-space representation $\left(\mathbb{C}^2,\ \alpha^2\right)$ of Γ^2_N. Especially, $\alpha^2\left(g\right)$ acting on $\mathbb{C}^2$ are in the sense of (2.1.8), for all $g \in \Gamma^2_N$. i.e.,*

(2.1.9) $\left(\mathbb{C}^2,\ \alpha^2\right)$ *forms a 2-dimensional representation of* Γ^2_N,

where α^2 is in the sense of (2.1.8).

Proof. Since α^2 of (2.1.8) is a generator-preserving group-isomorphism from Γ^2_N onto $\mathfrak{M}^2_2(N)$, it satisfies

$$\alpha^2\left(g_1g_2\right) = \alpha^2\left(g_1\right)\ \alpha^2\left(g_2\right) \text{ in } \mathfrak{M}^2_2(N),$$

for all $g_1,\ g_2 \in \Gamma^2_N$, and

$$\alpha^2\left(g^{-1}\right) = \left(\alpha^2\left(g\right)\right)^{-1} \text{ in } \mathfrak{M}_2^2(N),$$

for all $g \in \Gamma_N^2$.

Since $\mathfrak{M}_2^2(N) \subseteq M_2(\mathbb{C})$, the matrices $\alpha^2(g)$ are acting on $\mathbb{C}^2$, for all $g \in \Gamma_N^2$. Therefore, the pair $\left(\mathbb{C}^2,\ \alpha^2\right)$ forms a Hilbert-space representation of Γ_N^2. ∎

2.2. **Certain Order-2 Matrices in $M_n(\mathbb{C})$.** Now, let $n > 2$ in $\mathbb{N}$. And consider certain types of matrices in $M_n(\mathbb{C})$. In Section 2.1, we showed that our group Γ_N^2 of (1.0.1) is group-isomorphic to the multiplicative subgroup $\mathfrak{M}_2^2(N)$ of (2.1.5) in $M_2(\mathbb{C})$, by (2.1.7), and hence, we obtain a natural 2-dimensional representation $\left(\mathbb{C}^2,\ \alpha^2\right)$ of Γ_N^2 in (2.1.9). In this section, we construct a base-stone to extend the representation $\left(\mathbb{C}^2,\ \alpha^2\right)$ to arbitrary n-dimensional representations $(\mathbb{C}^n,\ \alpha^n)$ of Γ_N^2, for all $n \in \mathbb{N} \setminus \{1\}$.

To do that, we fix the matrices formed by

$$A_2(a,\, b) = \begin{pmatrix} a & b \\ \frac{1-a^2}{b} & -a \end{pmatrix},$$

as blocks of certain matrices in $M_n(\mathbb{C})$.

Assumption For our main purpose, we always assume that:

$$(a,\, b) \in \mathbb{C}_{-1}^{\times} \times \mathbb{C}^{\times},$$

whenever we have matrices $A_2(a,\, b)$ in the following text, as in (2.1.3)′, where

$$\mathbb{C}_{-1}^{\times} \overset{def}{=} \mathbb{C} \setminus \{0,\, 1\}.$$

□

Assume first that $n = 3$, and consider a matrix A_3 formed by

(2.2.1)

$$A_3 = \begin{pmatrix} 1 & c & \frac{-bc}{1-a} \\ 0 & a & b \\ 0 & \frac{1-a^2}{b} & -a \end{pmatrix} \in M_3(\mathbb{C}),$$

for some $a,\, b,\, c \in \mathbb{C}^{\times}$. As we have seen in (2.2.1), this matrix A_3 is regarded as the following block matrix,

(2.2.1)′

$$A_3 = \begin{pmatrix} (1) & \begin{pmatrix} c & \frac{-bc}{1-a} \end{pmatrix} \\ \begin{pmatrix} 0 \\ 0 \end{pmatrix} & A_2(a,\ b) \end{pmatrix} \text{ in } M_3(\mathbb{C}),$$

where $A_2(a,\, b)$ is in the sense of (2.1.1).

If we have (3×3)-matrix A in the sense of (2.2.1) understood as (2.2.1)′, then, from straightforward computation, one can obtain that

(2.2.2)

$$A_3^2 = I_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

the identity matrix of $M_3(\mathbb{C})$.

Let's define now a morphism

$$A_3 : \mathbb{C}_{-1}^{\times} \times \mathbb{C}^{\times} \times \mathbb{C}^{\times} \to M_3(\mathbb{C}) \text{ by}$$

(2.2.3)

$$A_3(a,\,b,\,c) = \begin{pmatrix} (1) & \begin{pmatrix} c & \frac{-bc}{1-a} \end{pmatrix} \\ \begin{pmatrix} 0 \\ 0 \end{pmatrix} & A_2\,(a,\ b) \end{pmatrix},$$

identified with the matrix (2.2.1) in $M_3(\mathbb{C})$, by (2.2.1)$'$.

By (2.2.2), for any $(a,\,b,\,c) \in \mathbb{C}^{\times}_{\sim 1} \times \mathbb{C}^{\times} \times \mathbb{C}^{\times}$, we have

$$\left(A_3(a,b,c)\right)^2 = I_3 \text{ in } M_3(\mathbb{C}).$$

Now, let $n = 4$ in $\mathbb{N}$, and consider a matrix A_4 formed by

(2.2.3)

$$A_4 = \begin{pmatrix} 1 & 0 & d & \frac{-bd}{1-a} \\ 0 & 1 & c & \frac{-bc}{1-a} \\ 0 & 0 & a & b \\ 0 & 0 & \frac{1-a^2}{b} & -a \end{pmatrix} \in M_4(\mathbb{C}),$$

for $c,\,d \in \mathbb{C}^{\times}$. Then, similarly, this matrix A_4 of (2.2.3) is regarded as the block matrix,

(2.2.4)

$$A_4 = \begin{pmatrix} (1) & \begin{pmatrix} 0 & d & \frac{-bd}{1-a} \end{pmatrix} \\ \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} & A_3(a,b,c) \end{pmatrix},$$

or

$$A_4 = \begin{pmatrix} I_2 & \begin{pmatrix} d & \frac{-bd}{1-a} \\ c & \frac{-bc}{1-a} \end{pmatrix} \\ O_{2,2} & A_2(a,\ b) \end{pmatrix}.$$

in $M_4(\mathbb{C})$, where $O_{2,2}$ is the (2×2)-*zero matrix* whose all entries are zeroes.

Then, by the direct computation, one obtains that

(2.2.5)

$$A_4^2 = I_4, \text{ the identity matrix of } M_4(\mathbb{C}).$$

So, similar to (2.2.3), we define a morphism

$$A_4 : \mathbb{C}^{\times}_{\sim 1} \times \left(\mathbb{C}^{\times}\right)^3 \to M_4(\mathbb{C}) \text{ by}$$

(2.2.6)

$$A_4(a,\,b,\,c,\,d) = \begin{pmatrix} I_2 & Q_{a,b}\,(c,d) \\ O_{2,2} & A_2(a,\ b) \end{pmatrix},$$

where I_2 is the (2×2)-identity matrix, $O_{2,2}$ is the (2×2)-zero matrix, and

$$Q_{a,b}(c,d) = \begin{pmatrix} d & \frac{-bd}{1-a} \\ c & \frac{-bc}{1-a} \end{pmatrix}.$$

The image $A_4(a,b,c,d)$ of (2.2.6) becomes a well-defined matrix in $M_4(\mathbb{C})$, by (2.2.3) and (2.2.4). Moreover, by (2.2.5), we have

(2.2.7)

$$(A_4(a,b,c,d))^2 = I_4 \text{ in } M_4(\mathbb{C}).$$

Let's consider one more step: let $n = 5$. Similarly, we define the following map,

$$A_5 : \mathbb{C}^\times_{-1} \times (\mathbb{C}^\times)^4 \to M_5(\mathbb{C}) \text{ by}$$

(2.2.8)

$$A_5\,(a_1,\ ...,\ a_5) = \begin{pmatrix} I_3 & Q_{a_1,a_2}(a_3,a_4,a_5) \\ O_{2,3} & A_2(a_1,\ a_2) \end{pmatrix},$$

in $M_5(\mathbb{C})$, for all $(a_1, ..., a_5) \in (\mathbb{C}^\times)^5$, where $A_2(a_1, a_2)$ is in the sense of (2.1.1), $O_{2,3}$ is the (2×3)-zero matrix, and I_3 is the (3×3)-identity matrix, and

$$Q_{a_1,a_2}(a_3,a_4,a_5) = \begin{pmatrix} a_5 & \frac{-a_2a_5}{1-a_1} \\ a_4 & \frac{-a_2a_4}{1-a_1} \\ a_3 & \frac{-a_2a_3}{1-a_1} \end{pmatrix}.$$

i.e.,

$$A_5(a_1, ..., a_5) = \begin{pmatrix} 1 & 0 & 0 & a_5 & \frac{-a_2a_5}{1-a_1} \\ 0 & 1 & 0 & a_4 & \frac{-a_2a_4}{1-a_1} \\ 0 & 0 & 1 & a_3 & \frac{-a_2a_3}{1-a_1} \\ 0 & 0 & 0 & a_1 & a_2 \\ 0 & 0 & 0 & \frac{1-a_1^2}{a_2} & -a_1 \end{pmatrix} \in M_5(\mathbb{C}).$$

From direct computation, one again obtain that

(2.2.9)

$$(A_5(a_1, ..., a_5))^2 = I_5, \text{ the identity matrix of } M_5(\mathbb{C}).$$

Inductively, for $n \geq 3$, we define the following map

$$A_n : \mathbb{C}^\times_{-1} \times (\mathbb{C}^\times)^{n-1} \to M_n(\mathbb{C}), \text{ by}$$

(2.2.10)

$$A_n\,(a_1,\ ...,\ a_n) = \begin{pmatrix} I_{n-2} & Q_{a_1,a_2}\,(a_3, ..., a_n) \\ O_{2,n-2} & A_2(a_1,\ a_2) \end{pmatrix},$$

for all $(a_1, ..., a_n) \in (\mathbb{C}^\times)^n$, where

$A_2(a_1, a_2)$ is in the sense of (2.1.1),
$O_{2,n-2}$ is the $(2 \times (n-2))$-zero matrix,
I_{n-2} is the $(n-2) \times (n-2)$ identity matrix,

and

$$Q_{a_1,a_2}(a_3,\ ...,\ a_n) = \begin{pmatrix} a_n & \frac{-a_2a_n}{1-a_1} \\ \vdots & \vdots \\ a_4 & \frac{-a_2a_3}{1-a_1} \\ a_3 & \frac{-a_2a_3}{1-a_1} \end{pmatrix}.$$

Then we obtain the following computation.

Theorem 2.4. *Let A_n be the morphism (2.2.10), and let $A_n(a_1, ..., a_n)$ be the image of A_n realized in the matricial algebra $M_n(\mathbb{C})$, for arbitrarily fixed $n \geq 3$ in $\mathbb{N}$. Then*

(2.2.11)

$$(A_n(a_1, ..., a_n))^2 = I_n,$$

the identity matrix of $M_n(\mathbb{C})$.

Proof. Let $n \geq 3$ be given in $\mathbb{N}$. Then, by (2.2.2), (2.2.7) and (2.2.9), we have

$$(A_k(a_1, ..., a_k))^2 = I_k \text{ in } M_k(\mathbb{C}),$$

for $k = 3, 4, 5$.

Now, without loss of generality take $n \geq 6$ in $\mathbb{N}$, generally. Then, by (2.2.10),

$$A_n(a_1, ..., a_n) = \begin{pmatrix} I_{n-2} & Q_{a_1,a_2}(a_3, ..., a_n) \\ O_{2,n-2} & A_2(a_1, a_2) \end{pmatrix}.$$

For convenience, we let

$$I \overset{denote}{=} I_{n-2}, \qquad Q \overset{denote}{=} Q_{a_1,a_2}(a_3, ..., a_n),$$

and

$$O \overset{denote}{=} O_{2,n-2}, \qquad A \overset{denote}{=} A_2(a_1, a_2).$$

i.e.,

$$A_n(a_1, ..., a_n) \overset{denote}{=} \begin{pmatrix} I & Q \\ O & A \end{pmatrix},$$

as a block matrix in $M_n(\mathbb{C})$. Then

$$(A_n(a_1, ..., a_n))^2 = \begin{pmatrix} I & Q \\ O & A \end{pmatrix} \begin{pmatrix} I & Q \\ O & A \end{pmatrix}$$

$$= \begin{pmatrix} I^2 + QO & IQ + QA \\ OI + AO & OQ + A^2 \end{pmatrix}$$

(2.2.12)

$$= \begin{pmatrix} I & IQ + QA \\ O & A^2 \end{pmatrix} = \begin{pmatrix} I & Q + QA \\ O & I_2 \end{pmatrix},$$

by (2.1.2).

So, to show (2.2.11), it is sufficient to show that

$$Q + QA = O_{n-2,2},$$

by (2.2.12).

Notice that

$$QA = \begin{pmatrix} a_n & \frac{-a_2a_n}{1-a_1} \\ \vdots & \vdots \\ a_4 & \frac{-a_2a_4}{1-a_1} \\ a_3 & \frac{-a_2a_3}{1-a_1} \end{pmatrix} \begin{pmatrix} a_1 & a_2 \\ \frac{1-a_1^2}{a_2} & -a_1 \end{pmatrix}$$

$$= \begin{pmatrix} a_1a_n + \frac{-a_2a_n(1-a_1^2)}{a_2(1-a_1)} & a_2a_n + \frac{a_1a_2a_n}{1-a_1} \\ \vdots & \vdots \\ a_1a_3 + \frac{-a_2a_3(1-a_1^2)}{a_2(1-a_1)} & a_2a_3 + \frac{a_1a_2a_3}{1-a_1} \end{pmatrix}$$

$$= \begin{pmatrix} -a_n & a_2a_n\left(1+\frac{a_1}{1-a_1}\right) \\ \vdots & \\ -a_3 & a_2a_3\left(1+\frac{a_1}{1-a_1}\right) \end{pmatrix}.$$

Thus,

$$Q + QA = \begin{pmatrix} a_n & \frac{-a_2a_n}{1-a_1} \\ \vdots & \vdots \\ a_3 & \frac{-a_2a_3}{1-a_1} \end{pmatrix} + \begin{pmatrix} -a_n & a_2a_n\left(1+\frac{a_1}{1-a_1}\right) \\ \vdots & \\ -a_3 & a_2a_3\left(1+\frac{a_1}{1-a_1}\right) \end{pmatrix}$$

$$= \begin{pmatrix} 0 & a_2a_n\left(\frac{-1}{1-a_1}+1+\frac{a_1}{1-a_1}\right) \\ \vdots & \vdots \\ 0 & a_2a_3\left(\frac{-1}{1-a_1}+1+\frac{a_1}{1-a_1}\right) \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{pmatrix}$$

$$= O_{n-2,2},$$

i.e.,

$$Q + QA = O_{n-2,2}.$$

So, for any $n \geq 6$,

(2.2.13)

$$(A_n(a_1,\ ...,\ a_n))^2 = \begin{pmatrix} I & Q+QA \\ O & I \end{pmatrix} = \begin{pmatrix} I & O_{n-2,2} \\ O & I \end{pmatrix} = I_n,$$

the identity matrix of $M_n(\mathbb{C})$.

Therefore, for any $n \geq 3$ in $\mathbb{N}$,

$$(A_n(a_1,\ ...,\ a_n))^2 = I_n \text{ in } M_n(\mathbb{C}).$$

∎

2.3. **n-Dimensional Hilbert-Space Representations of Γ_N^2.** Let $N \in \mathbb{N}$ be the fixed quantity, and fix $n \in \mathbb{N} \setminus \{1\}$. Also, let Γ_N^2 be our finitely presented group (1.0.1). For the fixed n, we take the n-tuples

(2.3.1)

$$W_k = (a_{k,1}, ..., a_{k,n}) \in \mathbb{C}_{-1}^{\times} \times (\mathbb{C}^{\times})^{n-1},$$

for $k = 1, ..., N$, and assume that $W_1, ..., W_N$ are "mutually distinct."

Observation and Notation As we assumed above, let $W_1, ..., W_N$ be mutually distinct in $\mathbb{C}_{-1}^{\times} \times (\mathbb{C}^{\times})^{n-1}$. For our purposes, one may further assume that these mutually distinct n-tuples satisfy

$$W_i \ncong W_j$$

in the sense that:

$$a_{i,l} \neq a_{j,l} \text{ in } \mathbb{C}^{\times}, \text{ for all } l = 1, ..., N.$$

If the n-tuples $W_1, ..., W_N$ satisfy the above stronger condition than the mutually-distinctness, we say they are *strongly mutually distinct.* □

Define the corresponding matrices,

(2.3.2)

$$X_k \overset{denote}{=} A_n(W_k) = \begin{pmatrix} I_{n-2} & Q_{a_{k,1},a_{k,2}}(a_{k,3}, ..., a_{k,n}) \\ O_{2,\,n-2} & A_2(a_{k,1},\ a_{k,2}) \end{pmatrix},$$

as in (2.2.10), where W_k are "strongly mutually distinct" n-tuples of (2.3.1), for all $k = 1, ..., N$.

Then, as in Section 2.1, one can get the multiplicative subgroup $\mathfrak{M}_n^2(N)$ of $M_n(\mathbb{C})$ by the (reduced) free group generated by $\{X_1, ..., X_N\}$. i.e.,

(2.3.3)

$$\mathfrak{M}_n^2(N) = \left\langle \{X_j\}_{j=1}^N \right\rangle \text{ in } M_n(\mathbb{C}),$$

where X_j's are in the sense of (2.3.2), for all $j = 1, ..., N$.

Definition 2.2. *We call the multiplicative subgroup $\mathfrak{M}_n^2(N)$ of $M_n(\mathbb{C})$, an n-dimensional, order-2, N-generator (sub-)group (of $M_n(\mathbb{C})$).*

Then, similar to the proof of the structure theorem (2.1.7), one can obtain the following generalized result.

Theorem 2.5. *Let $\mathfrak{M}_n^2(N)$ be an n-dimensional, order-2, N-generator group in $M_n(\mathbb{C})$. Then it is group-isomorphic to the group Γ_N^2 of (1.0.1). i.e.,*

(2.3.4)

$$\mathfrak{M}_n^2(N) \overset{Group}{=} \Gamma_N^2, \text{ for all } n \in \mathbb{N} \setminus \{1\}.$$

In particular, there exist generator-preserving group-isomorphisms $\alpha^n : \Gamma_N^2 \to \mathfrak{M}_n^2(N)$, such that

(2.3.5)

$$\alpha^n(x_j) = X_j, \text{ for all } j = 1, ..., N,$$

where x_j's are the generators of Γ_N^2, and X_j's are in the sense of (2.3.2).

Proof. Let $n = 2$. Then, by (2.1.7), the isomorphic relation (2.3.4) holds. Assume now that $n \geq 3$. Then the proof of (2.3.4) is similar to that of (2.1.7). Indeed, one can show that

$$\mathfrak{M}_n^2(N) \overset{\text{Group}}{=} \mathcal{F}_N \,/\, \mathcal{R}_N \overset{\text{Group}}{=} \Gamma_N^2,$$

for all $n \in \mathbb{N}$. Therefore, there exists a natural generator-preserving group-isomorphism α^n from Γ_N^2 onto $\mathfrak{M}_n^2(N)$ as in (2.3.5). ∎

The structure theorem (2.3.4) is the generalized result of (2.1.7). And the group-isomorphism α^n of (2.3.5) generalizes α^2 of (2.1.8). Therefore, we obtain the following theorem, generalizing (2.1.9).

Theorem 2.6. *Let Γ_N^2 be the group in the sense of (1.0.1), for some $N \in \mathbb{N}$. Then there exist n-dimensional Hilbert-space representation $(\mathbb{C}^n,\ \alpha^n)$ of Γ_N^2, where α^n are in the sense of (2.3.5), for all $n \in \mathbb{N} \setminus \{1\}$. i.e.,*

(2.3.6) $(\mathbb{C}^n,\ \alpha^n)$ *form n-dimensional representations of Γ_N^2,*

for all $n \in \mathbb{N}$.

Proof. If $n = 2$, as we have seen in (2.1.9), there exists a 2-dimensional Hilbert-space representation $\left(\mathbb{C}^2,\ \alpha^2\right)$, where α^2 is the group-isomorphism in the sense of (2.1.8).

Suppose $n \geq 3$ in $\mathbb{N}$. For such n, two groups Γ_N^2 and $\mathfrak{M}_n^2(N)$ are isomorphic with a group-isomorphism $\alpha^n : \Gamma_N^2 \to \mathfrak{M}_n^2(N)$ of (2.3.5), by (2.3.4). It shows that the images $\alpha^n(g)$ are $(n \times n)$-matrices acting on the n-dimensional Hilbert space $\mathbb{C}^n$. So, the pair $(\mathbb{C}^n,\ \alpha^n)$ forms an n-dimensional representation of Γ_N^2, for all $n \in \mathbb{N}$. ∎

The above two theorems show that, for a fixed group Γ_N^2 of (1.0.1), one has a system

$$\{(\mathbb{C}^n,\ \alpha^n)\}_{n=2}^{\infty}$$

of Hilbert-space representations, and the corresponding isomorphic groups

$$\left\{\mathfrak{M}_n^2(N)\right\}_{n=2}^{\infty},$$

acting on $\mathbb{C}^n$ (or, acting in $M_n(\mathbb{C})$).

Example 2.1. *Let Γ_2^2 be the finitely presented group,*

$$\Gamma_2^2 = \left\langle \{x_1,\ x_2\},\ \{x_1^2 = x_2^2\} \right\rangle.$$

Fix $n = 3$. Now, take the following strongly distinct triples,

$$W_1 = (t_1,\ t_2,\ t_3),\ W_2 = (s_1,\ s_2,\ s_3),$$

in $\mathbb{C}_{-1}^{\times} \times \left(\mathbb{C}^{\times}\right)^2$, and construct two matrices,

$$A_3(W_1) = \begin{pmatrix} 1 & t_3 & \frac{-t_2 t_3}{1-t_1} \\ 0 & t_1 & t_2 \\ 0 & \frac{1-t_1^2}{t_2} & -t_1 \end{pmatrix},\ A_3(W_2) = \begin{pmatrix} 1 & s_3 & \frac{-s_2 s_3}{1-s_1} \\ 0 & s_1 & s_2 \\ 0 & \frac{1-s_1^2}{s_2} & -s_1 \end{pmatrix},$$

in $M_3(\mathbb{C})$. The group $\mathfrak{M}_3^2(2)$ is established as the reduced free group

$$\langle \{A_3(W_1),\ A_3(W_2)\} \rangle$$

generated by $A_3(W_1)$ and $A_3(W_2)$ in $M_3(\mathbb{C})$. Then

$$\mathfrak{M}_3^2(2) \overset{Group}{=} \Gamma_2^2.$$

So, one has a natural 3-dimensional representation $\left(\mathbb{C}^3,\ \alpha^3\right)$, *where* α^3 *is the group-isomorphism satisfying*

$$\alpha^3\left(x_l\right) = A_3(W_l), \text{ for all } l = 1, 2.$$

3. Application: A Group Algebra $\mathcal{A}_2$ Induced by Γ_2^2

In Section 2, we showed that the finitely presented group Γ_N^2 of (1.0.1) can have its family of finite-dimensional Hilbert-space representations

$$\{(\mathbb{C}^n,\ \alpha^n)\}_{n=2}^{\infty},$$

since it is group-isomorphic to n-dimensional, order-2, N-generator groups $\mathfrak{M}_n^2(N)$ in $M_n(\mathbb{C})$, for all $n \in \mathbb{N} \setminus \{1\}$, where

$$\mathfrak{M}_n^2(N) = \left\langle \{A_n(W_k)\}_{k=1}^N \right\rangle \text{ in } M_n(\mathbb{C}),$$

where W_1, ..., W_N are strongly mutually distinct n-tuples of $\mathbb{C}_{\neq 1}^{\times} \times \left(\mathbb{C}^{\times}\right)^{n-1}$, for all $n \in \mathbb{N} \setminus \{1\}$.

3.1. **Algebraic Observation for Group Elements of Γ_2^2.** In this section, we concentrate on the case where $N = 2$, i.e., we study the group elements of Γ_2^2 in detail. As a special case of (2.3.4),

(3.1.1)

$$\left\langle \{x_1, x_2\},\ \{x_1^2 = x_2^2 = e_2\} \right\rangle = \Gamma_2^2 \overset{\text{Group}}{=} \mathfrak{M}_2^2(2).$$

Consider Γ_2^2 pure-algebraically. Each element g of Γ_2^2 has its expression,

(3.1.2)

$$g = x_{i_1}^{k_1} x_{i_2}^{k_2} \cdots x_{i_n}^{k_{i_n}}, \text{ for some } n \in \mathbb{N},$$

as in (1.0.3), for some $(i_1, ..., i_n) \in \{1, 2\}^n$, and $(k_1, ..., k_n) \in \mathbb{Z}^n$.

However, by the relation on Γ_2^2,

$$x_1^2 = e \overset{denote}{=} e_2 = x_2^2 \text{ in } \Gamma_2^2,$$

one has that

(3.1.3)

$$x_l^{-1} = x_l, \text{ for } l = 1, 2,$$

and

$$x_l^{2k+1} = x_l, \text{ for all } k \in \mathbb{N}.$$

In other words, the generators x_1 and x_2 are "self-invertible," and "indeed of order-2," in Γ_2^2. The above two conditions in (3.1.3) can be summarized by

(3.1.3)′

$$x_l^{2n+1} = x_l, \text{ for } l = 1, 2, \text{ for all } n \in \mathbb{Z}.$$

So, the general expression (3.1.2) of g is in fact

(3.1.4)

$$g = x_{j_1} x_{j_2} ... x_{j_n} \text{ in } \Gamma_2^2,$$

for some $(j_1, ..., j_n) \in \{1, 2\}^n$, for some $n \in \mathbb{N}$, by (3.1.3), or by (3.1.3)′.

By the characterization,

$$\Gamma_2^2 \overset{\text{Group}}{=} \mathcal{F}_2 \ / \ \mathcal{R}_2,$$

and by the definition of the noncommutative (non-reduced) free group $\mathcal{F}_2$, the expression (3.1.4) of g goes to

(3.1.5)

$$g = \begin{cases} e = e_2 & \text{or} \\ x_1 & \text{or} \\ x_2 & \text{or} \\ (x_1x_2)^n x_1 & \text{or} \\ (x_1x_2)^n & \text{or} \\ (x_2x_1)^n x_2 & \text{or} \\ (x_2x_1)^n, & \end{cases}$$

in Γ_2^2, for all $n \in \mathbb{N}$.

Proposition 3.1. *Let $g \in \Gamma_2^2$. Then g is only one of the forms in (3.1.5).* $\square$

Let $g \in \Gamma_2^2 \setminus \{e, x_1, x_2\}$. Say,

$$g = x_1x_2x_1x_2...x_1x_2x_1.$$

Then it is self-invertible. Indeed,

$$\begin{aligned} g^2 &= (x_1x_2...x_1x_2x_1)(x_1x_2...x_1x_2x_1) \\ &= x_1x_2...x_1x_2\left(x_1^2\right)x_2...x_1x_2x_1 \\ &= x_1x_2...x_1x_2ex_2...x_1x_2x_1 \\ &= x_1x_2...x_1\left(x_2^2\right)x_1...x_1x_2x_1 \\ &= ... \\ &= x_1^2 = e, \end{aligned}$$

i.e.,

(3.1.6)

$$((x_1x_2)^n x_1)^2 = e, \text{ for all } n \in \mathbb{N}.$$

Similarly, one obtains that

(3.1.7)

$$((x_2x_1)^n x_2)^2 = e \text{ in } \Gamma_2^2,$$

for all $n \in \mathbb{N}$.

Now, let

$$g = x_1x_2x_1x_2...x_1x_2 \text{ in } \Gamma_2^2.$$

Then

$$g^{-1} = x_2x_1x_2x_1...x_2x_1,$$

with

$$|g| = \left|g^{-1}\right| \text{ in } \mathbb{N},$$

where $|w|$ means the *length of* $w \in \Gamma_2^2$. (For example, if $g = x_1x_2x_1x_2x_1 \in \Gamma_2^2$, then $|g| = 5$.)

Indeed,

$$gg^{-1} = (x_1x_2...x_1x_2)(x_2x_1...x_2x_1) = e.$$

So, one has that

(3.1.8)

$$((x_1x_2)^n)^{-1} = (x_2x_1)^n,$$

and equivalently,

(3.1.9)

$$((x_2x_1)^n)^{-1} = (x_1x_2)^n,$$

for all $n \in \mathbb{N}$.

Proposition 3.2. *Let $g \in \Gamma_2^2 \setminus \{e\}$, and let $|g|$ means the length of g in $\{x_1, x_2\}$ in Γ_2^2.*

(3.1.10) If $|g|$ is odd in $\mathbb{N}$, then $g^{-1} = g$ in Γ_2^2.

(3.1.11) If $|g|$ is even in $\mathbb{N}$, then

$$g = (x_1x_2)^k, \text{ or } (x_2x_1)^k,$$

and

$$g^{-1} = (x_2x_1)^k, \text{ respectively, } (x_1x_2)^k,$$

for some $k \in \mathbb{N}$.

Proof. Suppose $|g|$ is odd in $\mathbb{N}$. Then, by (3.1.5), the group element g is one of

$$x_1, \text{ or } x_2, \text{ or } (x_1x_2)^n x_1, \text{ or } (x_2x_1)^n x_2,$$

for $n \in \mathbb{N}$. By (3.1.6) and (3.1.7), in such cases,

$$g^2 = e \text{ in } \Gamma_2^2.$$

So, the statement (3.1.10) holds.

Now, assume that $|g|$is even in $\mathbb{N}$. Then, by (3.1.5), this element g is either

$$(x_1x_2)^k, \text{ or } (x_2x_1)^k, \text{ for } k \in \mathbb{N}.$$

By (3.1.8) and (3.1.9), one has that

$$\Big((x_1x_2)^k\Big)^{-1} = (x_2x_1)^k,$$

respectively,

$$\big((x_2x_1)^k\big)^{-1} = (x_1x_2)^k,$$

for all $k \in \mathbb{N}$.

Therefore, the statement (3.1.11) holds, too. ∎

By the self-invertibility of the group-identity e, the generators x_1, x_2, and the group elements g with odd length $|g|$, we are interested in the cases where

$$|g| \text{ is even.}$$

Equivalently, we are interested in the cases where

$$g = (x_1x_2)^k, \text{ or } g = (x_2x_1)^k \text{ in } \Gamma_2^2.$$

3.2. **Analytic-and-Combinatorial Observation for Elements of Γ_2^2 in $M_2(\mathbb{C})$.** As we have seen, the group Γ_2^2 is group-isomorphic to the 2-dimensional, order-2, 2-generator subgroup $\mathfrak{M}_2^2(2)$ of $M_2(\mathbb{C})$ under our representation $\big(\mathbb{C}^2,\ \alpha^2\big)$. In particular, $\mathfrak{M}_2^2(2)$ is generated by two matrices

$$A_2(t_1,\ s_1), \text{ and } A_2(t_2,\ s_2)$$

for the strongly mutually distinct pairs $(t_1,\ s_1)$ and $(t_2,\ s_2)$ in $\mathbb{C}^\times \times \mathbb{C}^\times$. More precisely,

$$A_2(t_1,\ s_1) = \begin{pmatrix} t_1 & s_1 \\ \frac{1-t_1^2}{s_1} & -t_1 \end{pmatrix}, \text{ and } A_2(t_2,\ s_2) = \begin{pmatrix} t_2 & s_2 \\ \frac{1-t_2^2}{s_2} & -t_2 \end{pmatrix}$$

in $M_2(\mathbb{C})$.

Notation In the rest of this paper, we denote $A_2(t_1,\, s_1)$ and $A_2(t_2,\, s_2)$, by X_1 and X_2, respectively, for convenience. $\square$

As we mentioned at the end of Section 3.1, we are interested in group-elements g formed by

(3.2.1)

$$g = (x_1x_2)^k \in \Gamma_2^2, \text{ for } k \in \mathbb{N}.$$

i.e., under α^2, if g is not of the form of (3.2.1), we obtain a self-invertible matrix

$$\alpha^2(g), \text{ such that } \big(\alpha^2(g)\big)^2 = I_2$$

in $\mathfrak{M}_2^2(2) \subset M_2(\mathbb{C})$.

However, if g is of (3.2.1) in Γ_2^2, then

(3.2.2)

$$\alpha^2\left(g\right) = \alpha^2\left((x_1x_2)^k\right) = \left(X_1X_2\right)^k,$$

and hence,

$$\big(\alpha^2(g)\big)^{-1} = \alpha^2\left(g^{-1}\right) = \left(X_2X_1\right)^k,$$

in $\mathfrak{M}_2^2(2)$.

Observe that

(3.2.3)

$$(X_1X_2) + (X_2X_1) = \left(2t_1t_2 + \tfrac{s_1(1-t_2^2)}{s_2} + \tfrac{s_2(1-t_1^2)}{s_1}\right) I_2$$

from the straightforward computation in $M_2(\mathbb{C})$.

Consider now that

$$\det\left(X_1\right) = -1 = \det(X_2),$$

and

$$\det\left(X_1 + X_2\right) = \det\begin{pmatrix} t_1+t_2 & s_1+s_2 \\ \frac{1-t_1^2}{s_1} + \frac{1-t_2^2}{s_2} & -t_1-t_2 \end{pmatrix}$$

$$= -\left(t_1^2 + 2t_1t_2 + t_2^2\right) - \left(1 - t_1^2 + \tfrac{s_1\left(1-t_2^2\right)}{s_2} + \tfrac{s_2\left(1-t_1^2\right)}{s_1} + 1 - t_2^2\right)$$

$$= -t_1^2 - 2t_1t_2 - t_2^2 - 1 + t_1^2 - \tfrac{s_1\left(1-t_2^2\right)}{s_2} - \tfrac{s_2\left(1-t_1^2\right)}{s_1} - 1 + t_2^2$$

$$= -2t_1t_2 - \tfrac{s_1\left(1-t_2^2\right)}{s_2} - \tfrac{s_2\left(1-t_1^2\right)}{s_1} - 2,$$

i.e.

$$\det\left(X_1 + X_2\right) = -2 - \left(2t_1t_2 + \tfrac{s_1\left(1-t_2^2\right)}{s_2} + \tfrac{s_2\left(1-t_1^2\right)}{s_1}\right),$$

where $\det(\cdot)$ means the *determinant on* $M_2(\mathbb{C})$

So, one can get that

(3.2.4)

$$\det\left(X_1\right) + \det\left(X_2\right) - \det\left(X_1 + X_2\right) = 2t_1t_2 + \tfrac{s_1\left(1-t_2^2\right)}{s_2} + \tfrac{s_2\left(1-t_1^2\right)}{s_1}.$$

Lemma 3.3. *Let X_1 and X_2 be the generating matrices of $\mathfrak{M}_2^2(2)$ as above. Then there exists $\varepsilon_0 \in \mathbb{C}$, such that*

(3.2.5)

$$(X_1X_2) + (X_2X_1) = \varepsilon_0\ I_2 \text{ in } M_2(\mathbb{C}),$$

and

$$\begin{aligned}\varepsilon_0 &= 2t_1t_2 + \tfrac{s_1\left(1-t_2^2\right)}{s_2} + \tfrac{s_2\left(1-t_1^2\right)}{s_1}\\ &= \det(X_1) + \det(X_2) - \det(X_1+X_2).\end{aligned}$$

Proof. The proof of (3.2.5) is done by (3.2.3) and (3.2.4). ∎

Therefore, by (3.2.5), one can observe that

$$\begin{aligned}(X_1X_2 + X_2X_1)^2 &= (X_1X_2)^2 + X_1X_2X_2X_1\\ &\quad + X_2X_1X_1X_2 + (X_2X_1)^2\\ &= (X_1X_2)^2 + (X_2X_1)^2 + 2I_2,\end{aligned}$$

and hence,

$$\begin{aligned}(X_1X_2)^2 + (X_2X_1)^2 &= (X_1X_2 + X_2X_1)^2 - 2I_2\\ &= (\varepsilon_0 I_2)^2 - 2I_2 = \left(\varepsilon_0^2 - 2\right) I_2,\end{aligned}$$

where ε_0 is in the sense of (3.2.5). i.e., we obtain that

(3.2.6)

$$(X_1X_2)^2 + (X_2X_1)^2 = \left(\varepsilon_0^2 - 2\right) I_2.$$

By (3.2.5) and (3.2.6), we have that

$$(X_1X_2)^1 + (X_2X_1)^1 = \varepsilon_0 I_2,$$

and

$$(X_1X_2)^2 + (X_2X_1)^2 = \left(\varepsilon_0^2 - 2\right) I_2,$$

where

$$\varepsilon_0 = \det(X_1) + \det(X_2) - \det(X_1+X_2),$$

as in (3.2.5).

More generally, we obtain the following recurrence relation.

Theorem 3.4. *Let X_1 and X_2 be the generating matrices of $\mathfrak{M}_2^2(2)$ in $M_2(\mathbb{C})$. If we denote*

$$\mathfrak{X}_k \overset{denote}{=} (X_1X_2)^k + (X_2X_1)^k, \text{ for all } k \in \mathbb{N},$$

then the following recurrence relation is obtained:

(3.2.7)

$$\mathfrak{X}_1 = \varepsilon_0\ I_2, \text{ and } \mathfrak{X}_2 = \left(\varepsilon_0^2 - 2\right)\ I_2,$$

and

$$\mathfrak{X}_n = \varepsilon_0\ \mathfrak{X}_{n-1} - \mathfrak{X}_{n-2}, \text{ for all } n \geq 3 \text{ in } \mathbb{N},$$

in $M_2(\mathbb{C})$, where

$$\varepsilon_0 = \det(X_1) + \det(X_2) - \det(X_1+X_2).$$

Proof. By (3.2.5), indeed, one has

$$\mathfrak{X}_1 = \varepsilon_0\ I_2,$$

and by (3.2.6),

$$\mathfrak{X}_2 = \left(\varepsilon_0^2 - 2\right)\ I_2.$$

Suppose $n = 3$ in $\mathbb{N}$. Then

$$\begin{aligned}\mathfrak{X}_3 &= (X_1X_2)^3 + (X_2X_1)^3 \\ &= (X_1X_2 + X_2X_1)^3 - (3X_1X_2 + 3X_2X_1)\end{aligned}$$

by (3.1.5)

$$\begin{aligned}&= (\mathfrak{X}_1)^3 - 3\mathfrak{X}_1 = (\varepsilon_0 I_2)^3 - 3(\varepsilon_0 I_2) \\ &= \left(\varepsilon_0^3 - 3\varepsilon_0\right) I_2\end{aligned}$$

in $M_2(\mathbb{C})$.

Observe now that

$$\begin{aligned}\varepsilon_0\mathfrak{X}_2 - \mathfrak{X}_1 &= \varepsilon_0\left(\varepsilon_0^2 - 2\right) I_2 - \varepsilon_0 I_2 \\ &= \left(\varepsilon_0^3 - 2\varepsilon_0\right) I_2 - \varepsilon_0 I_2 \\ &= \left(\varepsilon_0^3 - 3\varepsilon_0\right) I_2,\end{aligned}$$

in $M_2(\mathbb{C})$.

Thus, one obtains that

(3.2.8)

$$\mathfrak{X}_3 = \left(\varepsilon_0^3 - 3\varepsilon_0\right) I_2 = \varepsilon_0\mathfrak{X}_2 - \mathfrak{X}_1.$$

So, if $n = 3$, then the relation (3.2.7) holds true.

Assume now that the statement

(3.2.9)

$$\mathfrak{X}_{n_0} = \varepsilon_0\mathfrak{X}_{n_0-1} - \mathfrak{X}_{n_0-2}$$

hold for a fixed $n_0 \geq 3$ in $\mathbb{N}$. Then

$$\begin{aligned}\mathfrak{X}_{n_0+1} &= \left((X_1X_2)^{n_0} + (X_2X_1)^{n_0}\right)(X_1X_2 + X_2X_1) \\ &\qquad - (X_1X_2)^{n_0}(X_2X_1) - (X_2X_1)^{n_0}(X_1X_2) \\ &= \left((X_1X_2)^{n_0} + (X_2X_1)^{n_0}\right)(\varepsilon_0 I) \\ &\qquad - \left((X_1X_2)^{n_0-1} + (X_2X_1)^{n_0-1}\right)\end{aligned}$$

by the assumption (3.2.9)

$$= \varepsilon_0\mathfrak{X}_{n_0} - \mathfrak{X}_{n_0-1},$$

i.e., for the fixed $n_0 \geq 3$,

(3.2.10)

$$\mathfrak{X}_{n_0+1} = \varepsilon_0\mathfrak{X}_{n_0} - \mathfrak{X}_{n_0-1}.$$

Since n_0 is arbitrary in $\mathbb{N} \setminus \{1, 2\}$, we can conclude that

$$\mathfrak{X}_{n+1} = \varepsilon\mathfrak{X}_n - \mathfrak{X}_{n-1},$$

for all $n \in \mathbb{N} \setminus \{1\}$, by the induction, because of (3.2.8), (3.2.9) and (3.2.10).

Therefore, we obtain the recurrence relation:

$$\mathfrak{X}_1 = \varepsilon_0 I_2,\ \mathfrak{X}_2 = \left(\varepsilon_0^2 - 2\right) I_2,$$

and

$$\mathfrak{X}_n = \varepsilon_0\mathfrak{X}_{n-1} - \mathfrak{X}_{n-2},$$

for all $n \geq 3$ in $\mathbb{N}$.

Equivalently, the relation (3.2.7) holds. ∎

By the recurrence relation (3.2.7), we obtain that:

$$\begin{aligned} &X_1X_2 + X_2X_1 = \varepsilon_0 I_2, \\ &(X_1X_2)^2 + (X_2X_1)^2 = \left(\varepsilon_0^2 - 2\right) I_2, \\ &(X_1X_2)^3 + (X_2X_1)^3 = \left(\varepsilon_0^3 - 3\varepsilon_0\right) I_2. \end{aligned}$$

and

$$\begin{aligned} (X_1X_2)^4 + (X_2X_1)^4 &= \varepsilon_0 \left(\varepsilon_0^3 - 3\varepsilon_0\right) I_2 - \left(\varepsilon_0^2 - 2\right) I_2 \\ &= \left(\varepsilon_0^4 - 4\varepsilon_0^2 + 2\right) I_2, \end{aligned}$$

in $M_2(\mathbb{C})$.

Inductively, one can verify that:

Corollary 3.5. *There exists a functional sequence $(f_n)_{n=1}^{\infty}$, such that (3.2.11)*

$$(X_1X_2)^n + (X_2X_1)^n = (f_n(\varepsilon_0))\, I_2,$$

in $M_2(\mathbb{C})$. □

For instance, if $(f_n)_{n=1}^{\infty}$ is in the sense of (3.2.11), then

$$f_1(z) = z,\ f_2(z) = z^2 - 2,\ f_3(z) = z^3 - 3z,$$

and

$$f_4(z) = z^4 - 4z^2 + 2,$$

etc.

The following theorem provides the refined result of (3.2.11).

Theorem 3.6. *Let X_1 and X_2 be the generating matrices of a 2-dimensional, order-2, 2-generator subgroup $\mathfrak{M}_2^2(2)$ of $M_2(\mathbb{C})$ in the sense of (3.2.1), and let*

$$\mathfrak{X}_l = (X_1X_2)^l + (X_2X_1)^l \in M_2(\mathbb{C}),$$

for all $l \in \mathbb{N}$. Then there exists a functional sequence $(f_n)_{n=1}^{\infty}$ such that

$$\mathfrak{X}_n = f_n(\varepsilon_0)\ I_2 \text{ in } M_2(\mathbb{C}),$$

where

$$\varepsilon_0 = \det(X_1) + \det(X_2) - \det(X_1 + X_2).$$

Moreover,

(3.2.12)

$$\begin{aligned} f_n(z) &= \sum_{k=0}^{[\frac{n}{2}]} (-1)^k \tfrac{n}{k} \begin{pmatrix} n-k-1 \\ k-1 \end{pmatrix} z^{n-2k} \\ &= \sum_{k=0}^{[\frac{n}{2}]} (-1)^k \tfrac{n}{n-k} \begin{pmatrix} n-k \\ k \end{pmatrix} z^{n-2k}, \end{aligned}$$

where

$$[\tfrac{n}{2}] = \textit{the maximal integer} \leq \tfrac{n}{2},$$

and

$$\begin{pmatrix} m \\ l \end{pmatrix} = \tfrac{m!}{l!(m-l)!}, \textit{ for all } m,\ l \in \mathbb{N},$$

for all $n \in \mathbb{N}$.

Proof. By (3.2.7) and (3.2.11), there exists a functional sequence $(f_n)_{n=1}^{\infty}$ such that

$$\mathfrak{X}_n = f_n(\varepsilon_0)\ I_2 \text{ in } M_2(\mathbb{C}), \text{ for all } n \in \mathbb{N},$$

where

$$\varepsilon_0 = \det(X_1) + \det(X_2) - \det(X_1 + X_2).$$

For instance,

(3.2.13)

$$f_1(z) = z,\ f_2(z) = z^2 - 2, \text{ and } f_3(z) = z^3 - 3z,$$

etc. So, it suffices to show that each n-th entry of the sequence $f_n(z)$ satisfies (3.2.12), for all $n \in \mathbb{N}$.

Say $n = 1$. Then the function $f_1(z)$ of (3.2.12) satisfies

$$f_1(z) = \sum_{k=0}^{[\frac{1}{2}]} (-1)^k \frac{n}{n-k} \begin{pmatrix} n-k-1 \\ k-1 \end{pmatrix} z^{n-2k}$$

$$= \sum_{k=0}^{0} (-1)^k \frac{n}{n-k} \begin{pmatrix} n-k \\ k \end{pmatrix} z^{n-2k}$$

$$= (-1)^0 \frac{1}{1-0} \begin{pmatrix} 1-0 \\ 0 \end{pmatrix} z^{1-0} = z,$$

satisfying

(3.2.14)

$$f_1(z) = z.$$

If $n = 2$, then the function $f_2(z)$ of (3.2.12) goes to

$$f_2(z) = \sum_{k=0}^{1} (-1)^k \frac{n}{n-k} \begin{pmatrix} n-k \\ k \end{pmatrix} z^{n-2k}$$

$$= (-1)^0 \frac{2}{2-0} \begin{pmatrix} 2-0 \\ 0 \end{pmatrix} z^{2-0} + (-1)^1 \frac{2}{2-1} \begin{pmatrix} 2-1 \\ 1 \end{pmatrix} z^{2-2}$$

$$= z^2 - 1,$$

satisfying

(3.2.15)

$$f_2(z) = z^2 - 1.$$

Therefore, the formula (3.2.12) holds true where $n = 1$, 2, by (3.2.13).

Now, we consider the functions $f_n(z)$ of (3.2.12) satisfy the recurrence relation;

(3.2.16)

$$f_1(z) = z,\ f_2(z) = z^2 - 1,$$

and

$$f_{n+1}(z) = z f_n(z) - f_{n-1}(z),$$

for all $n \geq 2$ in $\mathbb{N}$.

First of all, the functions f_1 and f_2 satisfy the initial condition of the relation (3.2.16), by (3.2.14) and (3.2.15). So, concentrate on the cases where $n \geq 2$ in $\mathbb{N}$.

Observe that

$$z f_n(z) - f_{n-1}(z)$$

$$= z\left(\sum_{k=0}^{[\frac{n}{2}]}(-1)^k\ \tfrac{n}{k}\begin{pmatrix} n-k-1 \\ k-1\end{pmatrix} z^{n-2k}\right)-\sum_{k=0}^{[\frac{n-1}{2}]}(-1)^k\tfrac{n-1}{k}\begin{pmatrix} n-k-2 \\ k-1\end{pmatrix} z^{n-1-2k}$$

$$= \sum_{k=0}^{[\frac{n}{2}]}(-1)^k\tfrac{n}{k}\begin{pmatrix} n-k-1 \\ k-1\end{pmatrix} z^{n-2k+1} - \sum_{k=1}^{[\frac{n+1}{2}]}(-1)^{k-1}\ \tfrac{n-1}{k-1}\begin{pmatrix} n-k-1 \\ k-2\end{pmatrix} z^{n-1-2k+2}$$

$$= \sum_{k=0}^{[\frac{n}{2}]}(-1)^k\tfrac{n}{k}\begin{pmatrix} n-k-1 \\ k-1\end{pmatrix} z^{n+1-2k}+\sum_{k=1}^{[\frac{n+1}{2}]}(-1)^k\tfrac{n-1}{k-1}\begin{pmatrix} n-k-1 \\ k-2\end{pmatrix} z^{n+1-2k}$$

$$= z^{n+1}+\sum_{k=1}^{[\frac{n}{2}]}(-1)^k\tfrac{n}{k}\begin{pmatrix} n-k-1 \\ k-1\end{pmatrix} z^{n+1-2k}+\sum_{k=1}^{[\frac{n+1}{2}]}(-1)^k\tfrac{n-1}{k-1}\begin{pmatrix} n-k-1 \\ k-2\end{pmatrix} z^{n+1-2k}$$

$$= z^{n+1} + \sum_{k=1}^{[\frac{n}{2}]}(-1)^k\left(\tfrac{n}{k}\begin{pmatrix} n-k-1 \\ k-1\end{pmatrix} + \tfrac{n-1}{k-1}\begin{pmatrix} n-k-1 \\ k-2\end{pmatrix}\right) z^{n+1-2k}$$

$$+\ (-1)^{[\frac{n+1}{2}]}\tfrac{n-1}{[\frac{n+1}{2}]-1}\begin{pmatrix} n-[\frac{n+1}{2}]-1 \\ [\frac{n+1}{2}]-1\end{pmatrix} z^{n+1-2[\frac{n+1}{2}]}$$

$$= z^{n+1}+\sum_{k=1}^{[\frac{n}{2}]}(-1)^k\left(\tfrac{n(n-k-1)!}{k(n-k-1-k+1)!(k-1)!} + \tfrac{(n-1)(n-k-1)!}{(k-1)(n-k+k+2)!(k-2)!}\right) z^{n+1-2k} + \mathfrak{Y}$$

where
(3.2.17)

$$\mathfrak{Y} = (-1)^{[\frac{n+1}{2}]}\tfrac{n-1}{[\frac{n+1}{2}]-1}\begin{pmatrix} n-[\frac{n+1}{2}]-1 \\ [\frac{n+1}{2}]-1\end{pmatrix} z^{n+1-2[\frac{n+1}{2}]},$$

and then

$$= z^{n+1} + \sum_{k=1}^{[\frac{n}{2}]}(-1)^k\left(\tfrac{n(n-k-1)!}{(n-2k)!k!} + \tfrac{n(n-k-1)!-(n-k-1)!}{(n-2k+1)!(k-1)!}\right) z^{n+1-2k} + \mathfrak{Y}$$

$$= z^{n+1} + \sum_{k=1}^{[\frac{n}{2}]}(-1)^k\left(\tfrac{(n-2k+1)n(n-k-1)!+kn(n-k-1)!-k(n-k-1)!}{k(n-2k+1)!(k-1)!}\right) z^{n+1-2k} + \mathfrak{Y}$$

$$= z^{n+1} + \sum_{k=1}^{[\frac{n}{2}]}(-1)^k\left(\tfrac{(n^2-2kn+n+nk-k)(n-k-1)!}{k(n-2k+1)!(k-1)!}\right) z^{n+1-2k} + \mathfrak{Y}$$

$$= z^{n+1} + \sum_{k=1}^{[\frac{n}{2}]}(-1)^k\left(\tfrac{(n^2-nk+n-k)(n-k-1)!}{k(n-2k+1)!(k-1)!}\right) z^{n+1-2k} + \mathfrak{Y}$$

$$= z^{n+1} + \sum_{k=1}^{[\frac{n}{2}]}(-1)^k\left(\tfrac{(n+1)(n-k)(n-k-1)!}{k(n-2k+1)!(k-1)!}\right) z^{n+1-2k} + \mathfrak{Y}$$

$$= z^{n+1} + \sum_{k=1}^{[\frac{n}{2}]}(-1)^k\ \tfrac{n+1}{k}\ \tfrac{(n-k)!}{(n-k-(k-1))!(k-1)!}\ z^{n+1-2k} + \mathfrak{Y}$$

$$= z^{n+1} + \sum_{k=1}^{[\frac{n}{2}]}(-1)^k\ \tfrac{n+1}{k}\begin{pmatrix} n-k \\ k-1\end{pmatrix} z^{n+1-2k} + \mathfrak{Y}$$

$$= \sum_{k=0}^{[\frac{n}{2}]} (-1)^k \ \tfrac{n+1}{k} \left(\begin{array}{c} n-k \\ k-1 \end{array} \right) z^{n+1-2k} + \mathfrak{Y},$$

i.e., for any $n \geq 2$, we obtain
(3.2.18)

$$zf_n(z) - f_{n-1}(z) = \sum_{k=0}^{[\frac{n}{2}]} (-1)^k \ \tfrac{n+1}{k} \left(\begin{array}{c} n-k \\ k-1 \end{array} \right) z^{n+1-2k} + \mathfrak{Y},$$

where $\mathfrak{Y}$ is in the sense of (3.2.17).

Notice that if $n \geq 2$ is even in $\mathbb{N}$, then

$$[\tfrac{n}{2}] = [\tfrac{n+1}{2}] \text{ in } \mathbb{N},$$

and hence, the first term of (3.2.18) contains the second one. Therefore, one has
(3.2.19)

$$n \in 2\mathbb{N} \Rightarrow zf_n(z) - f_{n-1}(z) = \sum_{k=0}^{[\frac{n+1}{2}]} (-1)^k \ \tfrac{n+1}{k} \left(\begin{array}{c} n-k \\ k-1 \end{array} \right) z^{n+1-2k},$$

if and only if

$$n \in 2\mathbb{N} \Longrightarrow zf_n(z) - f_{n-1}(z) = f_{n+1}(z).$$

Now, assume $n \geq 2$ is odd in $\mathbb{N}$. Then

$$\mathfrak{Y} = (-1)^{[\frac{n+1}{2}]} \left(\tfrac{2n-2}{n+1-2} \right) \left(\tfrac{\big(n-[\frac{n+1}{2}]-1\big)!}{\big(n-[\frac{n+1}{2}]-1-[\frac{n+1}{2}]+2\big)!\big([\frac{n+1}{2}]-2\big)!} \right) z^{n+1-2[\frac{n+1}{2}]}$$

$$= (-1)^{[\frac{n+1}{2}]} \left(\tfrac{2n-2}{n-1} \right) \left(\tfrac{\big(n-[\frac{n+1}{2}]-1\big)!}{\big(n-2[\frac{n+1}{2}]+1\big)!\big([\frac{n+1}{2}]-2\big)!} \right) z^0.$$

$$= (-1)^{[\frac{n+1}{2}]} \, (2) \left(\tfrac{\big(n-[\frac{n+1}{2}]\big)!}{\big(n-2[\frac{n+1}{2}]+1\big)!\big(n-[\frac{n+1}{2}]\big)\big([\frac{n+1}{2}]-2\big)!} \right) z^0.$$

$$= (-1)^{[\frac{n+1}{2}]} \left(\tfrac{n+1}{[\frac{n+1}{2}]} \right) \left(\tfrac{[\frac{n+1}{2}]-1}{n-[\frac{n+1}{2}]} \right) \left(\begin{array}{c} n-[\frac{n+1}{2}] \\ \\ [\frac{n+1}{2}]-1 \end{array} \right) z^0.$$

$$= (-1)^{[\frac{n+1}{2}]} \left(\tfrac{n+1}{[\frac{n+1}{2}]} \right) \left(\tfrac{n+1-2}{2n-n-1} \right) \left(\begin{array}{c} n-[\frac{n+1}{2}] \\ \\ [\frac{n+1}{2}]-1 \end{array} \right) z^0.$$

$$= \left((-1)^{[\frac{n+1}{2}]} \left(\tfrac{n+1}{[\frac{n+1}{2}]} \right) \left(\begin{array}{c} n-[\frac{n+1}{2}] \\ \\ [\frac{n+1}{2}]-1 \end{array} \right) \right) z^0.$$

i.e.,
(3.2.20)

$$n \in 2\mathbb{N}+1 \Longrightarrow \mathfrak{Y} = \left((-1)^{[\frac{n+1}{2}]} \left(\frac{n+1}{[\frac{n+1}{2}]} \right) \begin{pmatrix} n - [\frac{n+1}{2}] \\ \\ [\frac{n+1}{2}] - 1 \end{pmatrix} \right) z^0.$$

So, by (3.2.20), if $n \geq 2$ is odd in $\mathbb{N}$, then

(3.2.21)

$$zf_n(z) - f_{n-1}(z) = \sum_{k=0}^{[\frac{n+1}{2}]} (-1)^k \ \frac{n+1}{k} \begin{pmatrix} n-k \\ k-1 \end{pmatrix} z^{n+1-2k},$$

if and only if

$$zf_n(z) - f_{n-1}(z) = f_{n+1}(z).$$

Therefore, by (3.2.19) and (3.2.21), one can conclude that

(3.2.22)

$$n \geq 2 \text{ in } \mathbb{N} \Longrightarrow zf_n(z) - f_{n-1}(z) = f_{n+1}(z).$$

The above result (3.2.22) shows that

$$f_{n+1}(\varepsilon_0) = \varepsilon_0 f_n(\varepsilon_0) - f_{n-1}(\varepsilon_0), \text{ for } n \geq 2 \text{ in } \mathbb{N},$$

with

$$f_1(\varepsilon_0) = \varepsilon_0, \text{ and } f_2(\varepsilon_0) = \varepsilon_0^2 - 2.$$

Recall that, by (3.2.7),

$$\mathfrak{X}_1 = \varepsilon_0 \ I_2, \text{ and } \mathfrak{X}_2 = \left(\varepsilon_0^2 - 2\right) \ I_2,$$

and

$$\mathfrak{X}_{n+1} = \varepsilon_0 \mathfrak{X}_n - \mathfrak{X}_{n-1} = f_n(\varepsilon_0) \ I_2.$$

Therefore, one can conclude that if

$$f_n(z) = \sum_{k=0}^{[\frac{n}{2}]} (-1)^k \left(\frac{n}{k}\right) \begin{pmatrix} n-k-1 \\ k-1 \end{pmatrix} z^{n-2k},$$

then

$$\mathfrak{X}_n = f_n(\varepsilon_0) \ I_2, \text{ for all } n \in \mathbb{N}.$$

∎

By the recurrence relation (3.2.7), we obtain the above main theorem of Section 3.

Summary of Section 3.2 If X_1 and X_2 are the generating matrices of the 2-dimensional, order-2, 2-generator subgroup $\mathfrak{M}_2^2(2)$ of $M_2(\mathbb{C})$, then

(3.2.23)

$$(X_1X_2)^n + (X_2X_1)^n = f_n(\varepsilon_0) \ I_2,$$

in $M_2(\mathbb{C})$, where

$$f_n(z) = \sum_{k=0}^{[\frac{n}{2}]} (-1)^k \left(\frac{n}{k}\right) \begin{pmatrix} n-k-1 \\ k-1 \end{pmatrix} z^{n-2k},$$

and

$$\varepsilon_0 = \det(X_1) + \det(X_2) - \det(X_1 + X_2),$$

for all $n \in \mathbb{N}$. ■

Let x_1 and x_2 be the generators of the group Γ_2^2 of (1.0.1), and let

$$\alpha^2(x_1) = X_1 \text{ and } \alpha^2(x_2) = X_2$$

be the corresponding generating matrices of the isomorphic group $\mathfrak{M}_2^2(2)$ of Γ_2^2 in $M_2(\mathbb{C})$. The main result (3.2.23) shows that the generators x_1 and x_2 of Γ_2^2 induces the analytic data in $M_2(\mathbb{C})$ depending on the elements x_1x_2 and its inverses x_2x_1 up to (3.2.23).

3.3. **Group Algebras $A_{\alpha^n,\,\Gamma_N^2}$.** Let Γ be an arbitrary discrete group, and let $(H,\ \alpha)$ be a *Hilbert-space representation of* Γ consisting of a Hilbert space H and the group-action

$$\alpha : \Gamma \to L(H),$$

making

$$\alpha(g) : H \to H, \text{ for all } g \in \Gamma$$

be linear operators (or linear transformations) on H, satisfying

$$\alpha(g_1g_2) = \alpha(g_1)\ \alpha(g_2), \text{ for all } g_1,\ g_2 \in \Gamma,$$

and

$$\alpha(g^{-1}) = (\alpha(g))^{-1}, \text{ for all } g \in \Gamma,$$

where $L(H)$ means the operator algebra consisting of all linear operators on H.

Remark that, if we are working with *topologies*, then one may replace $L(H)$ to $B(H)$, the operator algebra consisting of all *bounded* (or *continuous*) linear operators on H. However, we are not considering topologies throughout this paper, so we simply set the representational models for $L(H)$ here.

For example, our group Γ_N^2 of (1.0.1) has its Hilbert-space representations $(\mathbb{C}^n,\ \alpha^n)$, for all $n \in \mathbb{N} \setminus \{1\}$, and each element g of Γ_N^2 is realized as $\alpha^n(g) \in \mathfrak{M}_n^2(N)$ in

(3.3.1)

$$M_n(\mathbb{C}) = L(\mathbb{C}^n) = B(\mathbb{C}^n).$$

The first equivalence relation (=) of (3.3.1) holds because, under finite-dimensionality, all linear operators on $\mathbb{C}^n$ are matrices acting on $\mathbb{C}^n$, and vice versa. The second equivalence relation (=) of (3.3.1) holds, because, under finite-dimensionality, all pure-algebraic operators on $\mathbb{C}^n$ are automatically bounded (and hence, continuous).

For an arbitrary group Γ, realized by $(H,\ \alpha)$, one obtains the isomorphic group $\alpha(\Gamma)$ in $L(H)$, i.e.,

$$\Gamma \overset{\text{Group}}{=} \alpha(\Gamma) \text{ in } L(H).$$

So, one can construct a subalgebra,

(3.3.2)

$$A_{\alpha,\,\Gamma} = \mathbb{C}\,[\alpha(\Gamma)] \text{ of } L(H),$$

where $\mathbb{C}[X]$ mean the *polynomial ring* in sets X. Such rings $\mathbb{C}[X]$ form algebras under polynomial addition and polynomial multiplication over $\mathbb{C}$, and hence, under the inherited operator addition and operator multiplication, $\mathbb{C}\,[\alpha(\Gamma)]$ forms an algebra in $L(H)$ in (3.3.2).

Definition 3.1. *Let Γ be a group and let $(H,\, \alpha)$ be a Hilbert-space representation of Γ. The subalgebra $A_{\alpha,\,\Gamma}$ of $L(H)$ in the sense of (3.3.2) is called the group algebra of Γ induced by $(H,\, \alpha)$.*

It is not difficult to show that if

(3.3.3)

$$A_\Gamma^o = \mathbb{C}[\Gamma],$$

the pure-algebraic algebra generated by Γ, as a polynomial ring in Γ, then

(3.3.4)

$$A_\Gamma^o \overset{\text{Alg}}{=} A_{\alpha,\,\Gamma},$$

for all Hilbert-space representations $(H,\, \alpha)$ of Γ algebraically, where "$\overset{\text{Alg}}{=}$" means "being algebra-isomorphic."

Proposition 3.7. *Let Γ_N^2 be the group of (1.0.1), and let $(\mathbb{C}^n,\, \alpha^n)$ be our n-dimensional representations of Γ_N^2, for all $n \in \mathbb{N} \setminus \{1\}$. Let $\mathfrak{M}_n^2(N)$ be the n-dimensional, order-2, N-generator subgroups of $M_n(\mathbb{C})$, for all $n \in \mathbb{N} \setminus \{1\}$. If $A_{\alpha^n,\,\Gamma_N^2}$ are in the sense of (3.3.2), and $A_{\Gamma_N^2}$ is the group algebra in the sense of (3.3.3), then*

(3.3.5)

$$A_{\Gamma_N^2} \overset{Alg}{=} A_{\alpha^n,\,\Gamma_N^2} = \mathbb{C}\left[\mathfrak{M}_n^2(N)\right], \text{ for all } n \in \mathbb{N}.$$

Proof. The proof of (3.3.5) is from the general relation (3.3.4). Recall that, by the group-isomorphism α^n,

$\alpha^n\left(\Gamma_2^2\right) = \mathfrak{M}_n^2(N)$, for all $n \in \mathbb{N}$.

∎

3.4. **A Group Algebra $A_{\alpha^2,\,\Gamma_2^2}$.** In this section, we take the group algebra in the sense of (3.3.2),

(3.4.1)

$$A_{\alpha^2,\Gamma_2^2} \overset{denote}{=} \mathcal{A}_2,$$

as a subalgebra of $M_2(\mathbb{C}) = L(\mathbb{C}^2)$, under our representation $\left(\mathbb{C}^2,\, \alpha^2\right)$. By (3.3.5), pure-algebraically, the algebra $\mathcal{A}_2$ of (3.4.1) is algebra-isomorphic to the group algebra $A_{\Gamma_2^2} = \mathbb{C}[\Gamma_2^2]$.

By (3.1.5), all elements of $\mathcal{A}_2$ are the linear combinations of the matrices formed by

(3.4.2)

$$I_2,\, X_1,\, X_2,\, (X_1X_2)^n,\, (X_2X_1)^n,$$

or

$$(X_1X_2)^n X_1,\, (X_2X_1)^n X_2,$$

for all $n \in \mathbb{N}$, where $X_j = \alpha^2(x_j)$, for $j = 1,\, 2$, since

$$\mathcal{A}_2 = \mathbb{C}\left[\mathfrak{M}_2^2(2)\right] \text{ in } M_2(\mathbb{C}).$$

As we have seen in Section 3.1, the building blocks of $\mathcal{A}_2$ in the sense of (3.4.2) satisfy that: (i) I_2, X_1, X_2, $(X_1X_2)^nX_1$, and $(X_2X_1)^nX_2$ are self-invertible in $\mathcal{A}_2$ $\subset M_2(\mathbb{C})$, and (ii)

$$((X_1X_2)^n)^{-1} = (X_2X_1)^n,$$

for all $n \in \mathbb{N}$.

Corollary 3.8. *Let X_1 and X_2 be the generating matrices of $\mathfrak{M}_2^2(2)$, and hence, those of $\mathcal{A}_2$. Then*

(3.4.3)

$$((X_1X_2)^n)^{-1} = f_n(\varepsilon_0)\ I_2 - (X_1X_2)^n,$$

and

$$((X_2X_1))^{-1} = f_n(\varepsilon_0)\ I_2 - (X_2X_1)^n$$

for all $n \in \mathbb{N}$, where $(f_n)_{n=2}^{\infty}$ is the functional sequence in the sense of (3.2.12).

Proof. By (3.2.23), we have

$$(X_1X_2)^n + (X_2X_1)^n = f_n(\varepsilon_0)\ I_2 \text{ in } \mathcal{A}_2,$$

where f_n are in the sense of (3.2.12), and ε_0 is in the sense of (3.2.5), for all $n \in \mathbb{N}$. So,

$$(X_2X_1)^n = f_n(\varepsilon_0)I_2 - (X_1X_2)^n \text{ in } \mathcal{A}_2,$$

for all $n \in \mathbb{N}$. Thus,

$$((X_1X_2)^n)^{-1} = f_n(\varepsilon_0)\ I_2 - (X_1X_2)^n,$$

for all $n \in \mathbb{N}$.
Similarly, we obtain

$$((X_2X_1)^n)^{-1} = f_n(\varepsilon_0)\ I_2 - (X_2X_1)^n,$$

for all $n \in \mathbb{N}$. ∎

It means that if T is a matrix of $\mathcal{A}_2$ in the form of either $(X_1X_2)^n$ or $(X_2X_1)^n$, for any $n \in \mathbb{N}$, then

$$T^{-1} = f_n(\varepsilon_0)\ I_2 - T.$$

Proposition 3.9. *Let $A \in \mathfrak{M}_2^2(2)$ in $\mathcal{A}_2$. Then*

(3.4.4)

$$A^{-1} = \begin{cases} A & \text{or} \\ r_A\ I_2 - A, & \end{cases}$$

for some $r_A \in \mathbb{C}$.

Proof. If $A \in \mathfrak{M}_2^2(2)$ in $\mathcal{A}_2$, then T is one of the forms of (3.4.2). As we discussed above, if A is I_2, or X_1, or X_2, or $(X_1X_2)^nX_1$, or $(X_2X_1)^nX_2$, for $n \in \mathbb{N}$, then it is self-invertible in $\mathcal{A}_2$. i.e.,

$$A^{-1} = A \text{ in } \mathcal{A}_2.$$

If A is either $(X_1X_2)^n$, or $(X_2X_1)^n$, for any $n \in \mathbb{N}$, then, by (3.4.3), there exists $r_A \in \mathbb{C}$, such that

$$A^{-1} = r_A - A \text{ in } \mathcal{A}_2 \vdots.$$

∎

The above proposition characterizes the invertibility of $\mathfrak{M}_2^2(2)$ "in $\mathcal{A}_2$."

3.5. **Trace of Certain Matrices of $\mathcal{A}_2$.** Throughout this section, we will use the same notations used before. Let $M_n(\mathbb{C})$ be the $(n \times n)$-matricial algebra, for $n \in \mathbb{N}$. Then the *trace tr on* $M_n(\mathbb{C})$ is well-defined as a linear functional on $M_n(\mathbb{C})$ by

$$tr\begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix} = \sum_{j=1}^{n} a_{jj}.$$

For instance, if $n = 2$, then

$$tr\begin{pmatrix} a & b \\ c & d \end{pmatrix} = a + d.$$

Now, let $\mathcal{A}_2$ be the group algebra of Γ_2^2 induced by $\left(\mathbb{C}^2,\ \alpha^2\right)$ in $M_2(\mathbb{C})$, in the sense of (3.4.1). Since the algebra $\mathcal{A}_2$ is a subalgebra of $M_2(\mathbb{C})$, one can naturally restrict the trace tr on $M_2(\mathbb{C})$ to that on $\mathcal{A}_2$. i.e., the pair $(\mathcal{A}_2,\ tr)$ forms a *noncommutative free probability space* in the sense of [8] and [9].

In this section, we are interested in trace of certain types of matrices in $\mathcal{A}_2$. Let X_1 and X_2 be the generating matrices of $\mathfrak{M}_2^2(2)$, and hence, those of $\mathcal{A}_2$. Define a new element T of $\mathcal{A}_2$ by

(3.5.1)

$$T = X_1 + X_2 \in \mathcal{A}_2.$$

By the self-invertibility of X_1 and X_2, this matrix is understood as the *radial operator* (e.g., [5]) *of* $\mathcal{A}_2$.

Observe that

$$\begin{aligned} T^2 &= (X_1 + X_2)^2 = X_1^2 + (X_1X_2 + X_2X_1) + X_2^2 \\ &= I_2 + f_1(\varepsilon_0)\ I_2 + I_2 = (f_1(\varepsilon_0) + 2)\ I_2, \end{aligned}$$

and

$$\begin{aligned} T^3 &= (X_1 + X_2)^3 = (X_1 + X_2)^2 (X_1 + X_2) \\ &= (f_1(\varepsilon_0) + 2)\, T, \end{aligned}$$

and

$$T^4 = T^3T = (f_1(\varepsilon_0) + 2)\, T^2 = (f_1(\varepsilon_0) + 2)^2\, I_2,$$

and

$$T^5 = T^4T = (f_1(\varepsilon_0) + 2)^2\, T,$$

and

$$T^6 = T^5T = (f_1(\varepsilon_0) + 2)^2\, T^2 = (f_1(\varepsilon_0) + 2)^3\, I_2,$$

etc, where f_n are in the sense of (3.2.12),

$$f_n(z) = \sum_{k=0}^{[\frac{n}{2}]} (-1)^k \left(\tfrac{n}{k}\right) \begin{pmatrix} n-k-1 \\ k-1 \end{pmatrix} z^{n-2k},$$

for all $n \in \mathbb{N}$, and hence,

$$f_1(\varepsilon_0) = \varepsilon_0.$$

Inductively, one obtains that:

Theorem 3.10. *Let $T = X_1 + X_2$ be the radial operator of $\mathcal{A}_2$, where X_1 and X_2 are the generating matrices of $\mathcal{A}_2$. Then*

(3.5.2)

$$T^{2n} = (\varepsilon_0 + 2)^n\, I_2, \text{ and } T^{2n+1} = (\varepsilon_0 + 2)^n\, T,$$

in $\mathcal{A}_2$, for all $n \in \mathbb{N}$, where

$$\varepsilon_0 = \det(X_1) + \det(X_2) - \det(X_1 + X_2).$$

Proof. The proof of (3.5.2) is done inductively by the observations in the very above paragraphs. Indeed, one can get that:

$$T^{2n} = (f_1(\varepsilon_0) + 2)^n I_2,$$

and

$$T^{2n+1} = (f_1(\varepsilon_0) + 2)^n T,$$

in $\mathcal{A}_2$, for all $n \in \mathbb{N}$. And, since $f_1(z) = z$, we obtain

$$f_1(\varepsilon_0) = \varepsilon_0,$$

where ε_0 is in the sense of (3.2.5). ∎

By (3.5.2), we obtain the following data.

Corollary 3.11. *Let $T = X_1 + X_2$ be the radial operator of $\mathcal{A}_2$. Then*
(3.5.3)

$$tr\left(T^n\right) = \begin{cases} 2\left(\varepsilon_0 + 2\right)^{n/2} & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd} \end{cases}$$

for all $n \in \mathbb{N}$.

Proof. Let's set
(3.5.4)

$$X_j = \begin{pmatrix} t_j & s_j \\ \frac{1-t_j^2}{s_j} & -t_j \end{pmatrix} \text{ in } \mathcal{A}_2, \text{ for } j = 1, 2,$$

where (t_1, s_1) and (t_2, s_2) are strongly distinct pair in $\mathbb{C}^\times \times \mathbb{C}^\times$.
By (3.5.2), if T is given as above in $\mathcal{A}_2$, then

$$T^{2n} = (\varepsilon_0 + 2)^n \ I_2, \text{ and } T^{2n+1} = (\varepsilon_0 + 2)^n T,$$

for all $n \in \mathbb{N}$.
Thus, it is not difficult to check that

$$tr\left(T^{2n}\right) = tr\left(\begin{pmatrix} (\varepsilon_0 + 2)^n & 0 \\ 0 & (\varepsilon_0 + 2)^n \end{pmatrix}\right)$$

$$= (\varepsilon_0 + 2)^n + (\varepsilon_0 + 2)^n = 2\left(\varepsilon_0 + 2\right)^n,$$

for all $n \in \mathbb{N}$, i.e.,
(3.5.5)

$$tr(T^{2n}) = 2\left(\varepsilon_0 + 2\right)^n, \text{ for all } n \in \mathbb{N}.$$

By the direct computation and by (3.5.4), one has
(3.5.6)

$$T = \begin{pmatrix} t_1 + t_2 & s_1 + s_2 \\ \frac{1-t_1^2}{s_1} + \frac{1-t_2^2}{s_2} & -(t_1 + t_2) \end{pmatrix}$$

in $\mathcal{A}_2$. So, one can get that:

$$tr(T) = (t_1 + t_2) + (-(t_1 + t_2)) = 0,$$

by (3.5.6).

Also, again by (3.5.2), we have

$$\begin{aligned} tr(T^{2n+1}) &= tr\left((\varepsilon_0 + 2)^n T\right) \\ &= (\varepsilon_0 + 2)^n \, tr(T) = 0, \end{aligned}$$

for all $n \in \mathbb{N}$. i.e.,

$$tr(T) = 0 = tr\left(T^{2n+1}\right), \text{ for all } n \in \mathbb{N}.$$

In summary, we have

(3.5.7)

$$tr\left(T^{2n-1}\right) = 0, \text{ for all } n \in \mathbb{N}.$$

Therefore, by (3.5.5) and (3.5.7), we obtain the formula (3.5.3). ∎

4. Application: $\mathcal{A}_2$ and Lucas Numbers

In this section, we will use the same notations used in Section 3 above. For instance, let

$$\mathcal{A}_2 = \mathbb{C}\left[\alpha^2(\Gamma_2^2)\right] = \mathbb{C}\left[\mathfrak{M}_2^2(2)\right]$$

is the group algebra of Γ_2^2 induced by our 2-dimensional representation $\left(\mathbb{C}^2,\ \alpha^2\right)$ in $M_2(\mathbb{C})$.

Here, as application of Sections 2 and 3, we study connections between analytic data obtained from elements of $\mathcal{A}_2$ and *Lucas numbers*.

For more details about Lucas numbers, see [7], [8], [9], [10], [11] and cited papers therein.

From the main result (3.2.23) of Section 3, we find a functional sequence $(f_n)_{n=1}^{\infty}$, with its n-th entries

(4.1)

$$f_n(z) = \sum_{k=0}^{[\frac{n}{2}]} (-1)^k \left(\tfrac{n}{k}\right) \begin{pmatrix} n-k-1 \\ k-1 \end{pmatrix} z^{n-2k}.$$

for all $n \in \mathbb{N}$, satisfying

$$(X_1X_2)^n + ((X_1X_2)^n)^{-1} = f_n(\varepsilon_0)\ I_2 \text{ in } \mathcal{A}_2,$$

where

$$\varepsilon_0 = \det(X_1) + \det(X_2) - \det\left(X_1 + X_2\right).$$

For any arbitrarily fixed $n \in \mathbb{N}$, consider the (non-alternative parts of) summands

(4.2)

$$\left(\tfrac{n}{k}\right) \begin{pmatrix} n-k-1 \\ k-1 \end{pmatrix}, \text{ for } k = 0,\ 1,\ ...,\ [\tfrac{n}{2}],$$

of $f_n(z)$ in (4.1).

Define such quantities of (4.2) by a form of a function,

$$g : \mathbb{N} \times \mathbb{N}_0 \to \mathbb{C} \text{ by}$$

(4.3)

$$g(n,\ k) \overset{def}{=} \begin{cases} \left(\frac{n}{k}\right) \begin{pmatrix} n-k-1 \\ k-1 \end{pmatrix} & \text{if } k = 0, 1,\ ..., [\frac{n}{2}] \\ 0 & \text{otherwise,} \end{cases}$$

for all $(n,\ k) \in \mathbb{N} \times \mathbb{N}_0$, where

$$\mathbb{N}_0 \overset{def}{=} \mathbb{N} \cup \{0\}.$$

Note that the definition (4.3) of the function g represents the (non-alternating parts) of summands of $(f_n)_{n=1}^{\infty}$.

The authors are not sure if the following theorem is already proven or not. They could not find the proofs in their references. So, we provide the following proof of the theorem.

Theorem 4.1. *Let g be a function from $\mathbb{N} \times \mathbb{N}_0$ into $\mathbb{C}$ be in the sense of (4.3), Then*

(4.4)

$$g(1,\ 0) = 1,$$

and

$$g\,(n+1,\ k) = g(n,\ k) + g(n-1,\ k-1),$$

for all $(n,\ k) \in (\mathbb{N} \setminus 1) \times \mathbb{N}$.

Proof. Observe first that:

(4.5)

$$g(n,k) \quad = \frac{n}{k} \begin{pmatrix} n-k-1 \\ k-1 \end{pmatrix} = \frac{n}{k} \frac{(n-k-1)!}{(k-1)!(n-k-1-k+1)!}$$

$$= \frac{n(n-k-1)!}{(n-2k)!k!},$$

whenever $g(n,\ k)$ is non-zero, for $(n,\ k) \in \mathbb{N} \times \mathbb{N}_0$.

By (4.5), we obtain that

(4.6)

$$g(1,\ 0) = \frac{1(1-0-1)!}{(1-0)!0!} = 1,$$

$$g(2,\ 0) = \frac{2(2-0-1)!}{(2-0)!0!} = 1,$$

and

$$g(2,\ 1) = \frac{2(2-1-1)!}{(2-2)!1!} = 2.$$

Consider now that:

$$g(n,\ k) + g(n-1,\ k-1)$$

$$= \frac{n}{k} \begin{pmatrix} n-k-1 \\ k-1 \end{pmatrix} + \frac{n-1}{k-1} \begin{pmatrix} n-1-(k-1)-1 \\ (k-1)-1 \end{pmatrix}$$

$$= \frac{n(n-k-1)!}{(n-2k)!k!} + \frac{(n-1)(n-k-1)!}{(n-2k+1)!(k-1)!}$$

by (4.5)

$$= \frac{n(n-2k+1)(n-k-1)!+k(n-1)(n-k-1)!}{(n-kk+1)!k!}$$

$$= \frac{(n^2+nk+n-k)(n-k-1)!}{(n-2k+1)!k!}$$

$$= \frac{(n+1)(n-k)!}{k(2n-2k+1)(k-1)!} = \frac{n+1}{k} \begin{pmatrix} (n+1)-k-1 \\ k-1 \end{pmatrix},$$

and hence,

(4.7)

$$\begin{aligned} g(n,\ k) + g(n-1, k-1) &= \frac{n+1}{k} \begin{pmatrix} (n+1)-k-1 \\ k-1 \end{pmatrix} \\ &= g(n+1,\ k). \end{aligned}$$

Therefore, one can get the recurrence relation (4.4), by (4.6) and (4.7). ∎

By (4.4), we can realize that the family

$$\{g(n,\, k) : (n,\, k) \in \mathbb{N} \times \mathbb{N}_0\}$$

satisfies the following diagram

(4.8)

$$\begin{array}{ccccccccccccccc}
 & & & & & & & & & & & & & & 2 \\
 & & & & & & & & & & & & 1 & & \\
 & & & & & & & & & & 1 & & 2 & & \\
 & & & & & & & & & 1 & & 3 & & & \\
 & & & & & & & & 1 & & 4 & & 2 & & \\
 & & & & & & & 1 & & 5 & & 5 & & & \\
 & & & & & & 1 & & 6 & & 9 & & 2 & & \\
 & & & & & 1 & & 7 & & 14 & & 7 & & & \\
 & & & & 1 & & 8 & & 20 & & 16 & & 2 & & \\
 & & & 1 & & 9 & & 27 & & 30 & & 9 & & & \\
 & & 1 & & 10 & & 35 & & 50 & & 25 & & 2 & & \\
 & & & & & & & & & & & & & & \\
 & & & & & & \vdots & & & & \vdots & & & &
\end{array}$$

where the rows of (4.8) represents $n \in \mathbb{N}$, and the columns of (4.8) represents $k \in \mathbb{N}_0$ of the quantities $g(n,\, k)$.

Observation The above diagram (4.8) is nothing but the *Lucas triangle* (e.g., [11]) induced by the *Lucas numbers* (e.g., [7], [8], [9] and [10]). □

By using the above family $\{g(n,\ k)\}_{(n,\, k)\in\mathbb{N}\times\mathbb{N}_0}$ of quantities obtained from (4.3), one can re-write the n-th entries f_n of (4.1) as follows;

(4.9)

$$f_n(z) = \sum_{k=0}^{[\frac{n}{2}]} (-1)^k\ (g(n,\ k))\, z^{n-2k},$$

for all $n \in \mathbb{N}$.

The new expression (4.9) shows that the coefficients of f_n is determined by the Lucas numbers alternatively, by (4.8).

Corollary 4.2. *The coefficients of the functions f_n of (4.1) are determined by the Lucas numbers in the Lucas triangle (4.8) alternatively.* $\square$

Moreover, we can verify that

$$\begin{aligned}
f_{n+1}(z) &= \sum_{k=0}^{[\frac{n+1}{2}]} (-1)^k \left(g(n+1,\ k)\right) z^{n+1-2k} \\
&= \sum_{k=0}^{[\frac{n+1}{2}]} (-1)^k \left(g(n,\ k) + g(n-1,\ k-1)\right) z^{n+1-2k} \\
&= \sum_{k=0}^{[\frac{n+1}{2}]} (-1)^k \left(g(n,\ k)\right)\ z^{n+1-2k} \\
&\qquad + \sum_{k=1}^{[\frac{n+1}{2}]} (-1)^k \left(g(n-1,\ k-1)\right) z^{n-2k} \\
&= z\left(\sum_{k=0}^{[\frac{n}{2}]} (-1)^k \left(g(n,\ k-1)\right) z^{n-2k}\right) \\
&\qquad - \sum_{k=0}^{[\frac{n-1}{2}]} (-1)^k \left(g(n-1,\ k-1)\right) z^{n-1-2k} \\
&= z f_n(z) - f_{n-1}(z).
\end{aligned}$$

i.e., from the recurrence relation (4.7), we can re-prove the recurrence relation (3.2.16).

Corollary 4.3. *Let*

$$f_n(z) = \sum_{k=0}^{[\frac{n}{2}]} (-1)^k \left(g(n,\ k)\right) z^{n-2k}$$

be in the sense of (4.9), where $\{g(n,\ k)\}_{(n,k)\in\mathbb{N}\times\mathbb{N}_0}$ are in the sense of (4.3). Then

$$f_1(z) = z \text{ and } f_2(z) = z^2 - 2,$$

and

$$f_{n+1}(z) = z f_n(z) - f_{n-1}(z),$$

for all $n \geq 2$ in $\mathbb{N}$. $\square$

Again, the above corollary demonstrates the connection between our Hilbert-space representations of Γ_2^2 and Lucas numbers.

References

[1] I. Cho, Free Probability on Hecke Algebras and Certain Group C^*-Algebras Induced by Hecke Algebras, Opuscula Math., (2015) To Appear.

[2] I. Cho, Representations and Corresponding Operators Induced by Hecke Algebras, DOI: 10.1007/s11785-014-0418-7, (2014) To Appear.

[3] R. Speicher, Combinatorial Theory of the Free Product with Amalgamation and Operator-Valued Free Probability Theory, Momoir, Ame. Math. Soc., 627, (1998)

[4] D. Voiculescu, K. J. Dykemma, and A. Nica, Free Random Variables, CRM Monograph Ser., vol.1, ISBN-13: 978-0821811405, (1992)

[5] I. Cho, and T. Gillespie, Free Probability on Hecke Algebras, DOI: 10.1007/s11785-014-0403-1, (2014) To Appear.

[6] I. Cho, The Moments of Certain Perturbed Operators of the Radial Operator in the Free Group Factors $L(F_N)$, J. Anal. Appl., vol. 5, no.3, (2007) 137 - 165.

[7] M. Asci, B. Cekim, and D. Tasci, Generating Matrices for Fibonaci, Lucas and Spectral Orthogonal Polynomials with Algorithms, Internat. J. Pure Appl. Math., 34, no. 2, (2007) 267 - 278.

[8] M. Cetin, M. Sezer, and C. Süler, Lucas Polynomial Approach for System of High-Order Linear Differential Equations and Residual Error Estimation, Math. Problems Eng., Article ID: 625984, (2015)

[9] A. Nalli, and P. Haukkanem, On Generalized Fibonacci and Lucas Polynomials, Chaos, Solitons & Fractals, 42, (2009) 3179 - 3186.

[10] J. Wang, Some New Results for the (p, q)-Fibonacci and Lucas Polynomicals, Adv. Diff. Equ., 64, (2014)

[11] A. T. Benjamin, The Lucas Triangle Recounted, available at
http://www.math.hmc.edu/~benjamin/papers/LucasTriangle.pdf.

Student Author

Ryan Golden is a senior at St. Ambrose University double majoring in Mathematics and Biology with minors in Computer Science and Chemistry. He plans to pursue a PhD in Applied Mathematics, specifically Mathematical and Computational Neuroscience. While his main interests are in the applications of mathematics, he is passionate about theoretical mathematics as well, and is indebted to Dr. Cho for providing the opportunity to work with and learn from him throughout his undergraduate career.

Press Summary In this paper, the authors studied matricial representations of certain finitely presented groups generated by N-generators of order-2. As an application, they considered corresponding group algebras under representation. Specifically, they characterized the inverses of all group elements in terms of matrices in the group algebra. From the study of this characterization, the authors realized there are close relations between the trace of the radial operator of the algebras and the Lucas numbers appearing in the Lucas triangle.

Saint Ambrose Univ., Dept. of Math. & Stat., 518 W. Locust St., Davenport, Iowa, 52803, U. S. A. /

E-mail address: goldenryanm@sau.edu / choilwoo@sau.edu